

# Non-union Clavicular Fractures Treated With Anatomical Locking Plate.

Sarabjeet Kohli<sup>1</sup>, Nilesh Vishwakarma<sup>2</sup>, Shaival Chauhan<sup>3</sup>, Kuldeep Salgotra<sup>4</sup>

<sup>1</sup>Associate Professor, Department of orthopaedics, MGM University of health sciences, Navi Mumbai.

<sup>2,3</sup>Assistant Professor, Department of orthopaedics, MGM University of health sciences, Navimumbai.

<sup>4</sup>Professor and Medical superintendent, Department of orthopaedics, MGM University of health sciences, Navi Mumbai.

Received: December 2016

Accepted: December 2016

**Copyright:** © the author(s), publisher. Annals of International Medical and Dental Research (AIMDR) is an Official Publication of "Society for Health Care & Research Development". It is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

## ABSTRACT

**Background:** The study aimed to analyze the outcomes of clavicle fractures in adults with established non union of mid third clavicle treated surgically and outcome of fracture union in terms of functionality of shoulder. **Methods:** 14 Non union clavicle fractures were treated in the Orthopedics Department at our tertiary institute. The average duration of surgery was 6 months from the date of injury. All the clavicle fracture non unions were operated using anatomical plate on superior side. None of the fractures had undergone any bone grafting. The shoulder function was evaluated using standard Constant Murley scoring system and American shoulder and elbow score (ASES). **Results:** Average time since injury was 11 months (range 6- 22 months). The majority 88% of patients had road traffic accidents. All 14 patients had optimum patients outcome scores with improvement in constant score from 46 pre-operatively to 89 postoperatively whereas ASES score improved from 46.6 pre-operatively to 94.0 postoperatively. The majority of non-union clavicle fractures were dissatisfied due to cosmetic, overhead-restricted endurance in shoulder activities and clavicular shortening. **Conclusion:** The operative treatment of non-union clavicle yields significantly improved outcomes with respect to shoulder function.

**Keywords:** Non-union Clavicle fractures.

## INTRODUCTION

Clavicle fractures form 2.6 % of all fractures<sup>[1-4]</sup> combined and 75 % of them are located at mid-shaft.<sup>[5,9]</sup> The non union rate in mid-shaft clavicle fractures is less than 1%.<sup>[12,13]</sup> Traditionally these fractures have been treated non-operatively, even when displaced.<sup>[6,7]</sup> The fractures often go into malunion and shortening which can affect the shoulder function significantly. The fractures can lead to non union which can be a disability in young active patients.<sup>[8]</sup>

### Name & Address of Corresponding Author

Dr. Nilesh Vishwakarma,  
Assistant Professor,  
Department of orthopaedics,  
MGM University of health sciences, Navi Mumbai

The aim of this study is to present the outcome of surgical treatment of mid-shaft non-union clavicular fracture in adults by open reduction and internal fixation with superior surface placement of locked reconstruction plate.

## MATERIALS AND METHODS

This retrospective study involved 14 patients with completely displaced mid-clavicular fractures who underwent open reduction and internal fixation with superior plating in our hospital between June 2012 and July 2016. The study includes all patients with non-union mid-clavicular fractures, whatever the type of fracture (transverse, oblique, or comminuted).

### Inclusion criteria:

- 1) Un-united fractures of midshaft clavicle.
- 2) Minimum 6 months since index injury.
- 3) Lateral and medial third clavicle fractures.
- 4) Consenting patients.

### Exclusion criteria:

- 1) Acute fractures.
- 2) Pathological fractures.
- 3) Previous surgeries.
- 4) Ipsilateral concomitant limb fractures.
- 5) Smokers and alcoholics.

All operations were performed under general anaesthesia with inter-scalene block in semi beach chair position with the head turned away from the side of the operation. The incision was made over the antero-superior aspect of the clavicle. The fracture was reduced and stabilized by an anatomical locking plate placed superiorly. Inter-fragmentary screws were inserted if necessary. All atrophic non unions had iliac crest bone grafting done. The limb was supported with an arm sling post operatively. The patient was advised to carry out gentle pendulum exercises only. The x-rays were advised immediately postoperatively and at 6 and 12 weeks unless delayed union was found. Union was considered to have occurred if clinically the fracture site was non-tender and, no abnormal movement was demonstrable, and radiologically when callus was visible.



(a)



(b)



(c)

**Figure 1: Clavicular plate fixation(a,b,c).**

The variables studied were mode of injury, injury-surgery interval, preoperative and postoperative Constant Murley<sup>[10]</sup> and American shoulder and elbow score<sup>[11]</sup> (ASES) along with and time required for union were recorded. The scores were calculated postoperatively at 6 months after surgery. The constant score<sup>[10]</sup> is a simple, 100-point score, and gives a reliable indication of shoulder function. It assesses pain (15), activities of daily living (20), range of movement (40) and power (25). The Constant score of the postoperative shoulder was compared with the preoperative shoulder at follow-up.

In ASES score<sup>[11]</sup>, the final pain score is calculated via an independent formula while the raw score from the functional questions is multiplied by a coefficient (to get the final score for the functional questions). The pain and functional portions are then summed to obtain the final ASES score with higher scores indicating better outcomes

## RESULTS

This study includes 14 patients, 32 male and 2 female, with the mean age of 31 years. Average time since injury was 11 months (range 6- 22 months). 22% of patients had road traffic accidents while 88 % had history of fall on outstretched hand . 6 clavicles were fractured on the right side and 8 on the left. 9 cases had atrophic non unions necessitating bone-grafting whereas 5 cases had hypertrophic non-union. All 14 patients had optimum patients outcome scores with improvement in constant score from 46 pre-operatively to 89 postoperatively whereas ASES score improved from 46.6 pre-operatively to 94.0 postoperatively

One patient developed an early wound infection, which was successfully managed by antibiotics. All patients were followed up until clinical and radiological union. Radiological union was defined as visible bridging callus or absence of a visible fracture line. The average time of union was 10 weeks (8-20 weeks). Union was delayed beyond 12 weeks in 4 cases. All the patients were relatively satisfied with the procedure. None of the patients had implant loosening or implant failure.



(a)



(b)



(c)

**Figure 2: Preoperative(a), immediate postoperative (b) and united fracture(c) x-rays.**

## DISCUSSION

Clavicle fractures form 2.6 % of all fractures combined and 75 % of them are located at mid-shaft.<sup>[9]</sup> The non union rate in mid-shaft clavicle fractures is less than 1%.<sup>[12,13]</sup> In contrast recent studies point at a non-union rate of 15% with patient outcomes turning out to be highly unsatisfactory (32%) as compared to studies like Altamimi et al<sup>[14]</sup> and Hill et al.<sup>[15]</sup> Neer<sup>[16]</sup> and Rowe<sup>[17]</sup> reported non union in less than 0.5% in their huge series with 2235 and 566 patients respectively. Malunion was also getting recognized more with its distinct cosmetic look. Nowak et al<sup>[18]</sup> studied complications in 208 adults with mid-clavicular fractures and found that, at ten years after the injury, ninety-six patients (46%) still had symptoms despite the fact that only fifteen (7%) had nonunion. This proved that patients can have major poor outcome following a clavicular mal-union because of symptoms specifically weakness and easy fatigability with overhead activities.

Mckee et al<sup>[19,20]</sup> showed clavicular shortening led to patient dissatisfaction. They noted poor abduction endurance. They concluded that some shortening with inferior displacement and anterior rotation of the lateral fragment seen in most cases. Thus shortening has a negative effect on muscle tendon unit.

Lazarides and Zafiropoulos et al<sup>[21]</sup> objectively showed that shortening of more than 18 mm in male patients and 14 mm in female patients was

associated with poor clinical outcome. Ledger et al<sup>[22]</sup>, 31 Eskola et al<sup>[23]</sup> Hill et al.<sup>[24]</sup> and Wick et al<sup>[25]</sup> also reported - poor clinical outcome if the shortening was more than 15 or 20 mm.

Most of the complications like brachial plexus palsy, vascular injury, Infection, plate failure, hypertrophic or dysesthetic scars, implant loosening or pneumothorax are rare in operative group.<sup>[26-28]</sup> Patients were more satisfied following operative intervention.<sup>[27]</sup> Correlation can exist between the degree of scapular mal-alignment due to clavicular shortening and shoulder dysfunction.<sup>[28-30]</sup>

The incidence of nonunion of the clavicle has been reported to be between 0.4 to 4.0%.<sup>[16,17]</sup> Nonunion is higher in the middle-third fractures of the clavicle followed by lateral-third fractures.<sup>[31-33]</sup> Nonunion is characterized by failure to show clinical or radiographic progression of healing at four to six months.<sup>[34-36]</sup> Nonunion may be either atrophic or hypertrophic, the former being a more common pattern than the latter.<sup>[37]</sup>

In our series of 14 cases of non union, there was one case of infection, excellent function with average constant murley score of 95.33 and all fractures had united in 20 weeks. None underwent reoperation for any complications. None had dissatisfaction except few (4 cases out of 14) who complained of prominent hardware not necessitating removal.

We conclude that superior plating of non united clavicular fractures heals optimally with higher patient satisfaction.

## CONCLUSION

Superior Anatomical plate fixation results in better outcomes and better patient satisfaction in non-union mid clavicular fractures.

## REFERENCES

1. Postacchini F, Gumina S, De Santis P, Albo P. Epidemiology of clavicle fractures. *J Shoulder Elbow Surg.* 2002; 11(5): 452-6.
2. CM Robinson. Fractures of the clavicle in the adult epidemiology and classification. *J Bone Joint Surg (Br).* 1998; 80: 476-84.
3. Preston CF, Egol KA. Midshaft Clavicle Fractures in Adults. *Bull NYU Hosp Jt Dis.* 2009; 67(1): 52-7.
4. Singh R, Rambani R, Kanakaris N and Giannoudis PV. A 2-year experience, management and outcome of 200 clavicle fractures. *Injury.* 2012; 43(2): 159-63.
5. Boehme D., Curtis R. J. Jr., De Haan J. T., et al. The treatment of non-union of fractures of the mid-shaft of the clavicle with an intramedullary Hagie pin and autogenous bone graft. *AAOS Instr. Course Lect.* 1993, 42, 283-290.
6. Rowe CR. An atlas of anatomy and treatment of midclavicular fractures. *Clin Orthop Relat Res.* 1968; 58: 29-42.
7. Sahal A. Altamimi: Nonoperative Treatment Compared with Plate Fixation of Displaced Midshaft Clavicular Fractures: Surgical Technique. *J Bone Joint Surg Am.* 2008; 90: 1-8.

8. Robinson, Court-Brown, McQueen, Wakefield. Estimating the risk of non-union following nonoperative treatment of a clavicular fracture. *J Bone Joint Surg Am.* 2004; 86: 1359-65.
9. Crenshaw AH. Fractures of the shoulder girdle arm and forearm. In: Crenshaw AH, editor. *Campbell's operative orthopaedics.* 8th ed. St.Louis: Mosby Year book; 1992.p 989-1053.
10. Constant CR, Murley AH. A clinical method of functional assessment of the shoulder. *Clin Orthop Relat Res.* 1987 Jan; (214):160-4.
11. American Shoulder and Elbow Surgeons Standardized Shoulder Assessment Form, patient self-report section: reliability, validity, and responsiveness. Michener LA, McClure PW, Sennett BJ. *J Shoulder Elbow Surg.* 2002 Nov-Dec; 11(6):587-94.
12. Eskola A, Vainionpaa S, Myllynen P, Patlala H, Rokkanen P. Outcome of clavicular fracture in 89 patients. *Arch Orthop Trauma Surg* 1986; 105: 337-8.
13. Stanley D. Recovery following fractures of the clavicle treated conservatively. *Injury* 1988; 19: 162-4.
14. Altamimi SA, McKee MD,. Nonoperative Treatment Compared with Plate Fixation of Displaced Midshaft Clavicular Fractures. *Surgical Technique. J Bone Joint Surg (Am).* 2008; 90(2); 1-8.
15. Hill JM, McGuire MH, Crosby LA. Closed treatment of displaced middle-third fractures of the clavicle gives poor results. *J Bone Joint Surg Br.* 1997; 79: 537-9.
16. Neer CS. Nonunion of the clavicle. *J Am Med Assoc* 1960;172: 1006-11.
17. Rowe CR. An atlas of anatomy and treatment of mid-clavicular fractures. *Clin Orthop.* 1968; 58: 29-42.
18. Nowak J, Holgersson M, Larsson S. Can we predict long-term sequelae after fractures of the clavicle based on initial findings? A prospective study with nine to ten years of follow-up. *J Shoulder Elbow Surg.* 2004;13:479-86.
19. McKee MD. Functional outcome following clavicle fractures in polytrauma patients. *J Trauma.* 1999; 47: 616.
20. McKee MD, Pedesen EM, Jones C, Stephen DJ, Kreder HJ, Schemitsch EH, Wild LM, Potter J. Deficits Following Nonoperative Treatment of Displaced Midshaft Clavicular Fractures. *J Bone Joint Surg (Am).* 2006; 88(1): 35-40.
21. Lazarides S, Zafiroopoulos G. Conservative treatment of fractures at the middle third of the clavicle: the relevance of shortening and clinical outcome. *J Shoulder Elbow Surg.* 2006; 15: 191-4.
22. Ledger M, Leeks N, Ackland T, Wang A. Short malunions of the clavicle: an anatomic and functional study. *J Shoulder Elbow Surg.* 2005; 14: 349-54.
23. Eskola A, Vainionpaa S, Myllynen P, Patlala H, Rokkanen P. Outcome of clavicular fracture in 89 patients. *Arch Orthop Trauma Surg.* 1986; 105: 337-8.
24. Hill JM, McGuire MH, Crosby LA. Closed treatment of displaced middle-third fractures of the clavicle gives poor results. *J Bone Joint Surg (Br).* 1997; 79: 537-9.
25. Wick M, Muller EJ, Kollig E, Muhr G. Midshaft fractures of the clavicle with a shortening of more than 2 cm predispose to nonunion. *Arch Orthop Trauma Surg* 2001; 121: 207-11.
26. Böstman O, Manninen M, Pihlajamäki H. Complications of plate fixation in fresh displaced midclavicular fractures. *J Trauma.* 1997; 43: 778-83.
27. Poigenfürst J, Rappold G, Fischer W. Plating of fresh clavicular fractures: results of 122 operations. *Injury.* 1992; 23: 237-41.
28. Freeland A. Unstable adult midclavicular fracture. *Orthopedics.* 1990; 13: 1279-81.
29. Rasmussen JV, Jensen SL, Petersen JB, Falstie-Jensen T, Lausten G and Olsen BS. A retrospective study of the association between shortening of the clavicle after fracture and the clinical outcome in 136 patients. *Injury.* 2011; 42(4): 414-7.
30. Ristevski B, Hall JA, Pearce D, Potter J, Farrugia M and McKee MD. The radiographic quantification of scapular malalignment after malunion of displaced clavicular shaft fractures. *J Shoulder Elbow Surg.* 2013; 22(2): 240-6.
31. Boyer M. I., Axelrod T. S. Atrophic nonunion of the clavicle : Treatment by compression plate, lag-screw fixation and bone graft. *J. Bone Joint Surg.* 1997; 79-B: 301-303.
32. Bradbury N., Hutchinson J., Hahn D., et al. Clavicular nonunion. 31/32 healed after plate fixation and bone grafting. *Acta Orthop. Scand.* 1996; 67: 367-370.
33. Davids P. H., Luitse J.S., Strating R.P., et al. Operative treatment for delayed union and nonunion of midshaft clavicular fractures AO reconstruction plate fixation and early mobilization. *J. Trauma.* 1996; 40: 985-986.
34. Ebraheim N. A., Mekhail A. O., Darwich M. Open reduction and internal fixation with bone grafting of clavicular nonunion. *J. Trauma.* 1997; 42: 701-704.
35. Olsen B. S., Vaesel M. T., Sojbjerg J. O. Treatment of midshaft clavicular nonunion with plate fixation and autologous bone grafting. *J. Shoulder Elbow Surg.* 1995; 4: 337- 344.
36. Wilkins R. M., Johnston R. M. Ununited fractures of the clavicle. *J. Bone Joint Surg.* 1983; 65-A: 773-778.
37. Pedersen M., Poulsen K. A., Thomsen F., et al. Operative treatment of clavicular nonunion. *Acta Orthop. Belg.* 1994; 60: 303-306

**How to cite this article:** Kohli S, Vishwakarma N, Chauhan S, Salgotra K. Non-union Clavicular Fractures Treated With Anatomical Locking Plate. *Ann. Int. Med. Den. Res.* 2017; 3(2):OR05-OR08.