

A Randomized Study Of Dexmedetomidine Used As Adjuvant In Epidural Post Operative Analgesia.

Kamala Kanta Pradhan¹, Ranjita Mohanty¹

¹Assistant Professor, Department of Anaesthesiology and Critical care, SCB Medical College. Cuttack, Odisha, India.

Received: October 2017

Accepted: October 2017

Copyright: © the author(s), publisher. Annals of International Medical and Dental Research (AIMDR) is an Official Publication of "Society for Health Care & Research Development". It is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ABSTRACT

Background: Epidural anaesthesia with adjuvants is commonly used for postoperative pain management in infraumbilical surgeries but search for ideal adjuvant without any side effect goes on. **Aim:** This study was done to evaluate the onset and duration of sensory and motor block and side effects of dexmedetomidine, when used as an adjuvant in epidural anaesthesia in infraumbilical surgeries. **Methods:** 60 patients of ASA status I and II, posted for infraumbilical surgeries were randomly allocated into two groups of 30 each. Group L patients received epidural 0.5% levobupivacaine. Group LD patients received epidural 0.5% levobupivacaine and 1µg/kg dexmedetomidine. Preoperative and postoperative block characteristics as well as hemodynamic parameters and side effects were monitored. **Results:** Dexmedetomidine as an adjuvant had an earlier onset and longer duration of sensory and motor block compared to levobupivacaine alone. Sedation scores were statistically significant with dexmedetomidine group in comparison to levobupivacaine alone group. **Conclusion:** Dexmedetomidine, as an adjuvant to epidural levobupivacaine provided prolonged analgesia in infraumbilical surgeries.

Keywords: Dexmedetomidine, levobupivacaine, infraumbilical surgeries

INTRODUCTION

Epidural anaesthesia provides both intra and post operative pain relief in various infraumbilical surgeries. Epidural bupivacaine had been commonly used in the past decades for providing adequate post-op pain relief.^[1] The analgesic duration can be prolonged by increasing dose of local anaesthetics; however the risk of systemic cardiac and neurotoxicity can be increased.^[2] Therefore, adjuvant can be added to local anaesthetics to prolong the analgesic duration so that the amount of local anaesthetics can be restricted. Recemic bupivacaine is most frequently used long acting local anaesthetic agent in regional anaesthesia. But the low dose bupivacaine is often used in order to reduce cardiovascular side effects which may not provide adequate postoperative analgesia.^[3] Levobupivacaine is the isolated S (-) isomer of bupivacaine. Due to lower affinity of S(-) isomer to cardiac sodium channel compared to R isomer, it is less cardio toxic.^[4] So we have chosen levobupivacaine as the local anaesthetic as it is longer acting and devoid of any cardiac side effects. Recently, several neuraxial adjuvants, including clonidine, opioids,

dexamethasone, ketamine, magnesium sulphate and midazolam have demonstrated the synergistic analgesic effect with local anaesthetics with varying degrees of success. But the search for ideal adjuvant for a particular local anaesthetic agent goes on.^[5] Literature is available using α -2 agonists like clonidine as adjuvant to local anaesthetics like bupivacaine, levobupivacaine and ropivacaine in epidural route but very few are there regarding their use with dexmedetomidine. α -2 adrenergic agonists like dexmedetomidine have both analgesic and sedative properties when used as an adjuvant in regional anaesthesia.^[6] Dexmedetomidine has an eight-fold greater affinity for α 2 adrenergic receptors than clonidine and much less α 1 activity. Its higher selectivity for α 2A receptors is responsible for the hypnotic and analgesic effects.^[7] Previous studies have shown that clonidine and dexmedetomidine improved the quality of block when used as adjuvant with ropivacaine or bupivacaine in epidural block but studies are limited where levobupivacaine is used with dexmedetomidine. This study was designed to compare the analgesic, sedative action and side effects of dexmedetomidine when added to levobupivacaine for epidural analgesia in patients undergoing infraumbilical surgeries.

Name & Address of Corresponding Author

Dr Ranjita Mohanty.
Assistant Professor
Department of Anaesthesiology and Critical care
SCB Medical College. Cuttack. Odisha. India

MATERIALS AND METHODS

Ethical committee approval and written informed consent were obtained from 60 ASA status (I / II)

patients of age 20-60 years, posted for infraumbilical surgeries. This study was done from Jan 2016 upto Jan 2017 in a tertiary care hospital. Patients with history of cardiac, respiratory, hepatic, neurological diseases and with allergy to the study drugs, were excluded from the study. ECG, pulse oximetry (SPO₂) and non-invasive blood pressure (NIBP) were monitored. Epidural anaesthesia was given in the sitting position, using 18 gauge tuohy epidural needle at L3-L4 space and epidural catheter was advanced 4 cm into the epidural space. A test dose of 3 ml of 2% lignocaine solution containing adrenaline 1: 200,000 was injected. After 4-6 min of injecting the test dose and excluding intravascular or subarachnoid injection study drug was injected. Patients were allocated into two groups in double blinded fashion. Group L received 17 ml levobupivacaine with 1ml normal saline and group LD received 17 ml of 0.5% levobupivacaine and 1µg/kg dexmedetomidine (diluted to 1ml) through epidural catheter. The study drugs were prepared by an anaesthetist who was blind about the study. Sensory block was assessed using the blunt end of a 27-gauge needle. Motor blockade was assessed by using the modified bromage scale (bromage 0: The patient is able to move the hip, knee and ankle; bromage 1: the patient is unable raise extended leg; bromage 2: The patient is unable to move the hip and knee but able to move the ankle; bromage 3: The patient is unable to move the hip, knee and ankle). The time to reach the peak sensory level and bromage 3 motor blocks were recorded before surgery. The regression time for sensory and motor block were recorded in post anaesthesia care unit. All durations were calculated from the time of epidural injection. The two groups were monitored pre and intraoperatively for heart rate, non-invasive blood pressure and O₂ saturation (SpO₂). Hypotension was defined as systolic blood pressure <90 mmHg or >30% decrease in baseline values and was treated by fluids and vasopressors. Tachycardia was defined as heart rate >100/min. Bradycardia was defined as heart rate <60/min and was treated by inj 0.6 mg atropine. Intraoperative nausea, vomiting, pruritus, sedation or any other side effects were recorded. Sedation was assessed by Ramsay sedation score (1: alert and awake, 2: arousable to verbal command, 3: arousable with gentle tactile stimulation, 4: arousable with vigorous shaking, 5: unarousable).^[8]

Statistical Methods

Data were presented as mean ± SD. t-test was used to compare the two groups for quantitative data and chi-square test was used for qualitative data by SPSS V18. Value of p<0.05 was considered statistically significant.

RESULTS

A total of 60 patients posted for infraumbilical surgeries were enrolled for the study. They were

randomly divided into two groups. The demographic profiles of the patients in both the groups were comparable with regards to age, sex, height, weight and body mass index. The ASA status of patients was similar in both the groups and mean duration of surgery was comparable in both the groups (p>0.05) [Table 1].

Table 1: Demographic profile of patients of both group.

| Demographic characteristics | L group (n=30) Mean ±SD | LD group (n=30) Mean ±SD | P value |
|--------------------------------|----------------------------|-----------------------------|---------|
| Age (yrs) | 48.2±11.5 | 47.2±10.8 | 0.34 |
| Sex (m:f) | 15:15 | 14:16 | 0.56 |
| Weight (kg) | 64.22±12.32 | 64.45±11.48 | 0.46 |
| Height (cm) | 154.8±9.58 | 153.35±9.8 | 0.34 |
| BMI(Kg/m ²) | 26.4±2.15 | 27.1±2.52 | 0.67 |
| ASA (I/II) | 28/2 | 27/3 | 1.0 |
| Mean duration of surgery (min) | 95.12±14.5 | 94.55±14.84 | 0.69 |

Onset of sensory block at T 10 level was earlier in group LD (6.98±2.51min) compared to the group L (8.34±2.95min). Higher dermatomal spread (T6-7) was seen in group LD in comparison to group L (T7-8). Time for maximum sensory level was shorter (11.32±4.46min) in group LD compared to group L (14.18±5.2min). All the above sensory block characteristics were statistically significant in group LD in comparison to group L except maximum sensory block level. Complete motor block was achieved earlier (14.75±6.24min) in group LD and 18.52±5.85min in group L which was statistically significant. (p<0.05). [Table 2].

Table 2: Comparison of preoperative block characteristics

| Block characteristics | L group (n=30) | LD group(n=30) | P Value |
|---|----------------|----------------|---------|
| Onset time of sensory block at T 10(mins) | 8.34±2.95 | 6.98±2.51 | 0.02 |
| Max sensory block level | T7-T8 | T6-T7 | 0.85 |
| Time to max sensory block (mins) | 14.18±5.2 | 11.32±4.46 | 0.001 |
| Time for complete motor block (mins) | 18.52±5.85 | 14.75±6.24 | 0.01 |
| Total ephedrine requirement (mg) | 7.90±2.5 | 5.95±2.8 | 0.12 |

Many previous studies had shown that dexmedetomidine can be used as intraoperative sedative agent. In our study mean sedation scores were significantly higher in group LD compared to group L which is statistically significant. [Table 3].

Table 3: Sedation score in both group

| Sedation score | L group(n=30) | LD group(n=30) | P Value |
|----------------|---------------|----------------|---------|
| 1 | 20 | 7 | 0.03 |
| 2 | 7 | 12 | 0.04 |
| 3 | 3 | 11 | 0.04 |
| 4 | 0 | 0 | |
| 5 | 0 | 0 | |

Mean time to 2 segmental dermatomal regression was 155.42 ± 9.54 min and 138.10 ± 8.15 min in group LD and group L respectively. Return of motor power to bromage 1 was 258.35 ± 27.82 min in L group and 285.27 ± 26.29 min in group LD. Both the block characteristics were statistically significant. The time for rescue analgesia was 325.94 ± 22.58 min in the group L and 354.58 ± 26.1 min in group LD which was statistically significant. ($P < 0.05$). [Table 4]. Parameters like heart rate, mean arterial pressure, spo2 and respiratory rate were stable and more or less similar in both the groups throughout the study period.

Table 4: Comparisons of post op block characteristics

| Post op block characteristics | L group (n=30) | LD group(n=30) | P Value |
|--|----------------|----------------|---------|
| Mean time to two segment regression (mins) | 138.10±8.15 | 155.42±9.54 | 0.0002 |
| Mean time to sensory regression at S1 (mins) | 295.35±32.24 | 342.12±33.15 | 0.0001 |
| Mean time to regression to bromage 1 (mins) | 258.35±27.82 | 285.27±26.29 | 0.0001 |
| Time to first rescue analgesia (mins) | 325.94±22.58 | 354.58±26.1 | 0.0001 |

Table 5: Comparison of side effects in intra and post-operative period.

| Side effect | Lgroup(n=30) | LD group(n=30) |
|------------------------|--------------|----------------|
| Nausea | 6 | 4 |
| Vomiting | 1 | 3 |
| Shivering | 4 | 3 |
| Headache | 0 | 2 |
| Dizziness | 0 | 0 |
| Dry mouth | 2 | 3 |
| Respiratory depression | 0 | 0 |

[Table 5] showed the comparative incidence of various side effects in both the groups which were statistically not significant. We did not observe respiratory depression in any patient in both the group.

DISCUSSION

Adjuvants are used with local anesthetics in the epidural anesthesia to prolong postoperative analgesia. Primary aim of these adjuvants is to fasten and prolong the sensory and motor block without any side effect. The pharmacologic properties of α -2 agonists like clonidine and dexmedetomidine have been used extensively in regional blocks. Epidural administration of these drugs is associated with sedation, analgesia, anxiolysis, hypnosis and sympatholysis.^[9] Clonidine has been used as adjuvant to local anesthetics successfully over the last few decade but it has its own side effects. Introduction of dexmedetomidine has raised question

about use of clonidine. The faster onset of action, and prolonged duration of analgesia in the post-operative period, makes it a very effective adjuvant to local anaesthetics in regional anaesthesia. In our study, dexmedetomidine was used as adjuvant to levobupivacaine in epidural anaesthesia in which levobupivacaine – dexmedetomidine combine produced earlier onset of epidural block, prolonged duration of sensory block and more sedation in comparison to levobupivacaine alone. There was no statistical difference in haemodynamic parameters in both groups. Disma et al in their study found that clonidine produced a local anaesthetic sparing effect with a dose dependent decrease in ED50 of levobupivacaine for caudal anaesthesia. In addition, there was a dose dependent prolongation of postoperative analgesia following lower abdominal surgery in children. A dose of $2 \mu\text{g kg}$ of clonidine provided the optimum balance between improved analgesia and minimal side effects.^[10] Wallet et al in their study found that the addition of clonidine to epidural levobupivacaine and sufentanil for patient controlled epidural analgesia in labour improved analgesia, reduced the supplementation rate and reduced pruritus.^[11] Milligan et al opined that, in patients undergoing total hip replacement, the addition of the alpha (2)-adrenergic agonist clonidine to epidural infusions of levobupivacaine significantly improved postoperative analgesia.^[12] Akin et al in their study found that caudal clonidine prolonged the duration of analgesia produced by caudal levobupivacaine without causing significant side effects.^[13] Mahran et al opined that both clonidine and fentanyl can be used as effective additive to epidural levobupivacaine for postoperative analgesia after radical cystectomy with no significant difference between them.^[14] Manal et al in a comparative study of epidural morphine and epidural dexmedetomidine used as adjuvant to levobupivacaine, found that dexmedetomidine was a good alternative to morphine as an adjuvant to levobupivacaine in epidural anaesthesia for major abdominal surgeries.^[15] Zeng XZ et al in their study found that low-dose epidural dexmedetomidine improved thoracic epidural anaesthesia for nephrectomy. Sensory and motor blockade duration was longer in the dexmedetomidine group than in the control group. Pain score and analgesic requirement was lower in dexmedetomidine group.^[16] Ahmed Sobhy Basuni et al used dexmedetomidine as adjuvant to low-dose levobupivacaine in spinal anaesthesia for knee arthroscopy. They opined that dexmedetomidine was a good alternative to fentanyl for supplementation with low-dose levobupivacaine.^[17] Aliye Esmaoglu et al concluded that intrathecal dexmedetomidine when added to levobupivacaine for spinal anaesthesia shortens sensory and motor block onset time and prolongs block duration without any significant adverse effects.^[18] Our study found

similar findings using dexmedetomidine as adjuvant to epidural levobupivacaine. El-Hennawy et al studied the effect by adding clonidine or dexmedetomidine to bupivacaine in caudal block in children. They found that addition of dexmedetomidine or clonidine to caudal bupivacaine significantly prolonged analgesia in children undergoing lower abdominal surgeries with no significant advantage of dexmedetomidine over clonidine.^[19] Al-Mustafa et al. used dexmedetomidine as an intrathecal adjuvant to bupivacaine and found that its use accelerated the onset of sensory block to reach T10 dermatome.^[20] Bajwa et al showed in their study that dexmedetomidine was a better adjuvant than clonidine in epidural ropivacaine anaesthesia in providing better intra-operative and postoperative analgesia.^[21] Wu H-H et al in a retrospective study opined that neuraxial dexmedetomidine was a favorable adjuvant to local anaesthetics which provides prolonged analgesia. Neuraxial dexmedetomidine was associated with good sedation scores and lower analgesic requirements and stable intra operative hemodynamic.^[22] All the above studies showed that dexmedetomidine was a potent adjuvant to levobupivacaine in epidural anaesthesia. It provided earlier onset and prolonged sensory block. Patient comfort, satisfaction and anxiolysis was better when dexmedetomidine was used as adjuvant to levobupivacaine in epidural anaesthesia.

CONCLUSION

Dexmedetomidine when used as an adjuvant to epidural levobupivacaine provided prolonged sensory and motor block without any significant side effects. Epidural dexmedetomidine with levobupivacaine provided prolonged analgesia and delayed the requirement of 1st rescue analgesia compared to levobupivacaine alone.

REFERENCES

1. Tziavrangos E, Schug SA. Regional anaesthesia and perioperative outcome. *Current opinion in anaesthesiology* 2006 ;19:521-525
2. Becker DE, Reed KL. Local anesthetics: review of pharmacological considerations. *Anesthesia progress* 2012;59:90-101.
3. Ulke ZS, Senturk M. Non-analgesic effects of thoracic epidural anesthesia. *Agriculture* 2007;19:6-12.
4. Huang Y, Pryor M, Mather L, Veering B. Cardiovascular and central nervous system effects of intravenous levobupivacaine and bupivacaine in sheep. *Anesth Analg* 1998;86:797-804.
5. Bardsley H, Gristwood R, Baker H, et al. A comparison of the cardiovascular effects of levobupivacaine and rac-bupivacaine following intravenous administration to healthy volunteers. *Brint J. Clin Pharmacol* 1998; 46:245-9.
6. Kamibayashi T, Maze M. Clinical uses of alpha-2 adrenergic agonists. *Anesthesiology* 2000;93:1345-9.
7. Scafati A. Analgesia and alpha agonists 2 *Medens Rev* 2004;4:7.

8. Mauro VA, Brandao ST. Clonidine and dexmedetomidine through epidural route for postoperative analgesia and sedation in a cholecystectomy. *Rev bras Anesthesiol* 2004;4:1-10.
9. Kanazi GE, Aouad MT, Jabbour Khoury SL, et al. Effect of low dose dexmedetomidine or clonidine on the characteristics of bupivacaine spinal block. *Acta Anesthesiol scand* 2006;50:222-7.
10. Disma N, Frawley G, Mameli L. Effect of epidural clonidine on minimum local anesthetic concentration (ED50) of levobupivacaine for caudal block in children. *Paediatr Anaesth*. 2011 Feb;21 (2): 128-35.
11. Wallet F, Clement JH, Bouret C. Effects of a continuous low-dose clonidine epidural regimen on pain, satisfaction and adverse events during labour: a randomized, double-blind, placebo-controlled trial. *Eur J Anaesthesiol*. 2010 May; 27(5): 441-7.
12. Milligan KR, Convery PN, Weir P. The efficacy and safety of epidural infusions of levobupivacaine with and without clonidine for postoperative pain relief in patients undergoing total hip replacement. *Anesth Analg*. 2000 Aug; 91(2):393-7.
13. Akin A, Ocalan S, Esmaglu A. The effects of caudal or intravenous clonidine on postoperative analgesia produced by caudal levobupivacaine in children. *Pediatric anesthesia* 2010;20:350-355.
14. Mahran E, Ibrahim W. Is the combination of epidural clonidine- levobupivacaine has same analgesic efficacy and safety as the combination Fentanyl – levobupivacaine after radical cystectomy. *Egyptian journal of anaesthesia* 2014 April; 30(2): 143-147.
15. Manal M, Kamal, Sahar M. Talaat Comparative study of epidural morphine and epidural dexmedetomidine used as adjuvant to levobupivacaine in major abdominal surgery *Egyptian Journal of Anaesthesia*. 2014 April; 30 (2):137-141.
16. Zeng XZ; Xu YM, CuiXG. Low-dose epidural dexmedetomidine improves thoracic epidural anaesthesia for nephrectomy *Anaesthesia And Intensive Care* 2014 Mar.; 42(2): 185-90.
17. Ahmed Sobhy Basuni, Hoda Alsaied Ahmed Ezz. Dexmedetomidine as supplement to low-dose levobupivacaine in spinal anaesthesia for knee arthroscopy *Egyptian Journal of Anaesthesia* 2014 April;30(2): 149-153.
18. Aliye Esmaglu, Sumeysra Turk ,Adnan Bayram, Aynur Akin.. The effects of Dexmedetomidine Added to Spinal Levobupivacaine for Transurethral Endoscopic Surgery, *Balkan Med J* 2013; 30: 186-90.
19. M. E-Hennawy, A. M. Abd-Elwahab, A.M. Abd-Elmaksoud., Addition of clonidine or dexmedetomidine to bupivacaine prolongs caudal analgesia in children. *Br. J. Anaesth*. 2009;103 (2):268-274.
20. Al-Mustafa MM, Abu-Halaweh SA, Aloweida AS, et al. Effect of dexmedetomidine added to spinal bupivacaine for urological procedures. *Saudi Med J*. 2009;30:365-70.
21. Bajwa SJS. Bajwa SK, Kaur J, et al Dexmedetomidine and clonidine in epidural anaesthesia: a comparative evaluation. *Indian J Anaesthe* 2011; 55 (2): 116-21.
22. Wu H-H, Wang H-T, Jin J-J, et al. Does Dexmedetomidine as a Neuraxial Adjuvant Facilitate Better Anesthesia and Analgesia ? A Systematic Review and Meta-Analysis. *PLoS ONE* 2014 Mar 9(3):e93114.

How to cite this article: Pradhan KK, Mohanty R. A Randomized Study Of Dexmedetomidine Used As Adjuvant In Epidural Post Operative Analgesia. *Ann. Int. Med. Den. Res.* 2017; 3(6):AN07-AN10.

Source of Support: Nil, **Conflict of Interest:** None declared