

# Calcium, Zinc and Copper Levels in Clavicles of Human Foetuses.

Farah Ghaus<sup>1</sup>, Nusra Rahman<sup>2</sup>, Nafis Ahmad Faruqi<sup>3</sup>

<sup>1</sup>Associate Professor, Department of Anatomy, J N Medical College, Aligarh Muslim University, Aligarh-202002, India

<sup>2</sup>Senior Resident, Department of Anatomy, J N Medical College, Aligarh Muslim University, Aligarh-202002, India

<sup>3</sup>Professor, Department of Anatomy, J N Medical College, Aligarh Muslim University, Aligarh-202002, India.

Received: November 2017

Accepted: December 2017

**Copyright:** © the author(s), publisher. Annals of International Medical and Dental Research (AIMDR) is an Official Publication of “Society for Health Care & Research Development”. It is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

## ABSTRACT

**Background:** Calcium, zinc and copper are essential minerals for growth. **Methods:** Twenty five human foetuses were divided into five groups i.e. group I (<17 wks), group II (17-20 wks), group III (21-25 wks), group IV (26-30 wks), group V (> 30 wks) having 5 foetuses each. **Results:** Fifty clavicles obtained after dissection were used to determine concentration of aforementioned minerals in each clavicle separately. Except a significant rise of calcium level in group II clavicles compared to group I ones, its deposition kept pace with matrix formation in clavicles of rest of the groups. **Conclusion:** Uniform concentration of zinc and copper and their constant ratio throughout in developing clavicles were unique observations.

**Keywords:** Calcium, Clavicle, Copper, Zinc.

## INTRODUCTION

Minerals play immense role in normal foetal development.<sup>[1]</sup> Calcium is the most abundant mineral of foetal body store. Previous literature regarding bodily calcium are based on studies in lower mammals.<sup>[2-4]</sup> Most of human studies on bone calcium contents are based on information in postnatal life.<sup>[5-7]</sup> Its concentration in human foetal tissue will have merit if considered in different stages of development. Copper and zinc are trace elements,<sup>[1,8,9]</sup> if deficient can lead to reduced embryonic development.<sup>[1,10,11]</sup> Data regarding concentrations of aforementioned elements in tissues of developing foetuses are based on studies on lower mammals.<sup>[12,13]</sup> Givens and Macy gave a detailed account of total calcium in developing human foetal bodies.<sup>[14]</sup> Tabrizi and Pakdel reported levels of Ca,<sup>[15]</sup> Cu and Zn in sacrum of women during three trimesters and their newborns. Such studies highlighted the nutritional status of foetus instead of developmental aspects. Scanty reports are available showing the levels of minerals in particular bone emphasizing their concentration

in different stages of developing human foetuses. Clavicle being an atypical long bone was chosen for this purpose.<sup>[15]</sup> Previous literature has measured only calcium in developing human bone.<sup>[16]</sup> We have taken into consideration copper and zinc also in addition to calcium to further enhance the scope of study.

## MATERIALS AND METHODS

Twenty five human foetuses were collected from museum of Department of Anatomy, Jawaharlal Nehru Medical College, Aligarh, U.P. Research works on these foetuses are already in progress for which permission was granted by Ethical Committee of Institution to post graduate students of Department of Anatomy. The foetuses were divided into five groups [Table 1] i.e. group I (<17 wks), group II (17-20 wks), group III (21-25 wks), group IV (26-30 wks), group V (> 30 wks) having 5 foetuses each. Clavicles were dissected out and disarticulated to remove them, obtaining ten clavicles from each group. Each clavicle was cut into three pieces to determine three minerals i.e. calcium, copper and zinc. Each clavicular piece was weighed and dissolved in 5ml concentrated nitric acid. Solutions thus obtained were filtered through Whatmann number I filter paper. Solutions were used to measure calcium by the method of Henry and Dryer 17 and zinc and copper by Atomic Absorption Spectrophotometry.<sup>[18,19]</sup> Data

### Name & Address of Corresponding Author

Dr. Ghaus Farah  
Associate Professor,  
Department of Anatomy,  
J.N. Medical College, A.M.U, Aligarh, 202002,  
U.P., India.

were analyzed by using one way ANOVA followed by post hoc Bonferroni test.

## RESULTS

[Table 2] shows detailed clavicular mineral levels in different groups. Levels of calcium in clavicle in group I was 1.38 mg/g of bone while in groups II, III, IV and V, these were 2.8, 3.12, 2.87 and 3.13 mg/g respectively. There was no statistically significant difference among last four groups. Most interestingly, the calcium concentration was significantly low in group I foetuses. Values of zinc ranged from 1.55ppm to 2.15 ppm and those of copper from 0.17 ppm to 0.20 ppm but none showed statistically significant variation when compared with rest of the groups.

**Table 1: Subgrouping of human fetuses.**

Groups	Age (wks)	Number of clavicles		
		Right	Left	Total
I	<17 wks	5	5	10
II	17-20	5	5	10
III	21-25	5	5	10
IV	26-30	5	5	10
V	>30	5	5	10
Total no. of clavicles		25	25	50

**Table 2: Comparison of levels of calcium (mg/g dry bone), zinc (ppm) and copper (ppm) in developing human foetal clavicles between five groups of gestational ages.**

Minerals	Group I (n=10) Mean ± SD	Group II (n=10) Mean ± SD	Group III (n=10) Mean ± SD	Group IV (n=10) Mean ± SD	Group V (n=10) Mean ± SD
Calcium	# f % ^ 1.38 ± 0.17	*2.80 ± 0.65	*3.12 ± 0.38	*2.87 ± 0.96	*3.13 ± 0.26
Zinc	1.55 ± 0.58	1.94 ± 0.27	2.15 ± 0.36	1.94 ± 0.27	2.15 ± 0.36
Copper	0.17 ± 0.02	0.20 ± 0.03	0.19 ± 0.04	0.20 ± 0.03	0.19 ± 0.04

Data were presented as mean ± standard deviation. Analysis was done using one – way ANOVA followed by post hoc Bonferroni test. The \* depicts comparison with Group I, # depicts comparison with Group II, f depicts comparison with Group III, % depicts comparison with Group IV, ^ depicts comparison with Group V \*# / f / % / ^ = < 0.05 (Significant)

## DISCUSSION & CONCLUSION

To provide strength to bone, both inorganic component i.e. calcium and organic matrix matter. The former makes it hard and rigid while latter imparts resilience.<sup>[20]</sup> Clavicle is first bone to ossify and ossification centre can be seen as early 6 wks of intrauterine life.<sup>[21]</sup> This is responsible for rise of calcium level with great rate making the reading 2.8 mg/g in group II compared to 1.38 mg/g in group I. It is interesting to note that the calcium concentration in clavicles of rest of groups showed no change. This clearly indicates that calcium

deposition kept pace with matrix formation to maintain a constant strength in rest of groups.

High concentration of zinc in foetal bone compared to soft tissue is clear indicator of its importance in growth of hard tissue of body.<sup>[22]</sup> Zinc is necessary part of hormone systems, and a deficiency of this vital trace elements will lead to impaired growth.<sup>[23]</sup> Association of copper with zinc for normal bodily growth is a well known fact.<sup>[24]</sup> The ratio of copper to zinc is clinically more important than concentration of either of these trace elements.<sup>[24]</sup> One of the most common trace metal imbalances is elevated copper and depressed zinc. A higher level of zinc relative to copper in our study conforms with previous findings of zinc concentration in different foetal tissues. A uniform concentration of both zinc and copper throughout foetal life in developing clavicles supports the importance of zinc- copper ratio in growth process. Further studies on growing bones in experimental animals after artificially altering the concentration of minerals are needed to reach some conclusion regarding their significance.

## Acknowledgements

We are thankful to Dr Arsalan Moinuddin, Assistant Professor, Department of Physiology, Shridev Suman Subharti Medical College, Dehradun for his help in statistical analysis, Prof. Haroon Subhan Khan, Department of Medicine and Dr Syed Shariq Naeem, Assistant Professor, Department of Pharmacology, JNMC, AMU, Aligarh for their help in calcium, zinc and copper estimation.

## REFERENCES

- Underwood EJ. Trace Elements in Human and Animal Nutrition. 4th ed. Academic Press, New York. 1977 (<http://www.elsevier.com/books/trace-elements-in-human-and-animals-nutrition/underwood>)
- Chan EL and Swaminathan R. Calcium metabolism and bone calcium content in normal and oophorectomized rats consuming various levels of saline for 12 months. J Nutr 1998; 128(3): 633-639.
- Feaster JP, Hansard SL, Outler JC and Davis GK. Placental transfer of calcium in the rat. J Nutrition 1956; 58:399-406.
- Graham RW and Scothorne RJ. Calcium homeostasis in the foetal guinea pig. Q J Exp Physiol 1970; 55:44-53.
- Goret- Nicaise M and Dhem A. Comparison of the calcium content of different tissues present in the human mandible. Acta Anatomica 1985; 124:167-172.
- Tzaphildou M and Zaichick V. Calcium, phosphorus, calcium phosphorus ratio in rib bone of healthy humans. Biol Trace Elem Res 2003; 93:63-74.
- Fischer A, Wiechula D, Postek- Stefan'ska L. and Kwapulin'ski J. Concentrations of metals in maxilla and mandible deciduous and permanent human teeth. Biol Trace Elem Res 2009 Dec; 132(1-3):19-26.
- Mertz W. Trace elements in Human and Animal Nutrition. 5th ed. Academic Press, San Diego, CA. 1987 (<https://www.elsevier.com/books/trace-elements-in-human-and-animal-nutrition/mertz/978-0-08-092468-7>).

9. Shaw JCL. Trace elements in the fetus and young infant II. Copper, Manganese, Selenium, and Chromium. Am J Dis Child 1980; 134(1):74-81.
10. Dreosit IE ,PC Grey, PJ Williams . Deoxyribosenucleic acid synthesis, protein synthesis and teratogenesis in zinc deficient rats. S Afr Med J 1972; 46:1585-1588.
11. Shaw JCL . Trace elements in the fetus and young infant I. Zinc .Am J Dis Child 1979 ; 133(12):1260-1268.
12. Abdelrahman MM. Copper, Manganese, Zinc, Iron and Calcium in fetal Tissue of Baladi Goats at Northern of Jordan. J of Animal and Vet Ad 2003; 2(4):209-214.
13. Bellof G, Most E, Pallauf J. Concentration of copper, iron, manganese and zinc in muscle, fat and bone tissue of lambs of the breed German Merino Landsheep in the course of the growing period and different feeding intensities. J. Anim Physiol Anim Nutr (Berl) 2007; 91(3-4):100-8.
14. Givens MH and Macy IG . The chemical composition of the human fetus. J Biol Chem 1933; 102:7-17.
15. Tabrizi FM and Pakdel FG. Serum level of some minerals during three trimesters of pregnancy in Iranian women and their newborns: A longitudinal study. Indian J Clin Biochem 2014; 29(2):174-180.
16. Ghaus F,Faruqi NA, Khan HS and Kirmani F. Calcium levels in the maxillae of human foetuses. Int.J. Morphol 2011; 29(1): 268-271.
17. Henry RJ and Dryer RL. In D. SELIGSON,Standard Methods of Clinical Chemistry.Vol 4, Academic Press, New York.1963; p 205-237.
18. Butrimovitz GP and Purdy WC. The Determination of Zinc in Blood Plasma By Atomic Absorption Spectrometry. Anal Chim Acta 1977 ; 94: 63-73.
19. Dawson JB, Ellis DJ and Newton-John H, Direct Estimation of Copper In Serum and Urine By Atomic Absorption Spectroscopy. Clin Chim Acta 1968; 21: 33.
20. Soames RW Skeletal system. In:Williams PM, editor. Gray's Anatomy.38th ed. Churchill Livingstone, 2000, p 425-736.
21. Faruqi N.A. The Clavicle, In: Human Osteology (A Clinical Orientation). 2nd ed. CBS Publishers, 2007 ,p 12-17.
22. Bergman Bo. Concentration of Zinc in some hard and soft tissues of rat determined by neutron activation analysis.Acta Radiologica: Therapy Physics Biology 1970; 9(5):430-432.
23. Khoshabi F,Shadan MR,Miri A and Rad JS. Determination of maternal serum, zinc, iron, calcium and magnesium during pregnancy in pregnant women and umbilical cord blood and their association with outcome of pregnancy. Mater Sociomed 2016; 28(2):104-107.
24. Osredkar J and Sustar N .Copper and Zinc, Biological role and significance of copper/zinc imbalance. J. Clinic Toxicol 2011; S3:001.

**How to cite this article:** Ghaus F, Rahman N, Faruqi NA. Calcium, Zinc and Copper Levels in Clavicles of Human Foetuses. Ann. Int. Med. Den. Res. 2018; 4(2):AT01-AT03.

**Source of Support:** Nil, **Conflict of Interest:** None declared