

Body Mass Index and its Correlation with Knee Pain in Osteoarthritis Patients.

Deepak Kumar Saxena¹, Atul Panday¹

¹Assistant Professor, Department of Orthopaedics, RMRI, Bareilly, Uttar Pradesh, India.

Received: May 2018

Accepted: June 2018

Copyright: © the author(s), publisher. Annals of International Medical and Dental Research (AIMDR) is an Official Publication of "Society for Health Care & Research Development". It is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ABSTRACT

Background: The goal of this study was to evaluate the correlation between body mass index (BMI) and severity of knee pain. **Methods:** Total 105 OA patients with knee pain selected on the basis of radiological diagnosis of OA knee were included. Evaluation of OA was done by using Kallgren and Lawrence grading along with numerical pain rating scale (NPRS). **Results:** Findings of the present study showed that number of OA patients 12 (11.4%), 40(38.09%), 30 (28.57%) and 33 (31.4%) were suffering with grade 1 to 4 OA severities respectively. Further, Pearson correlation coefficient showed a positive correlation between BMI and NPRS ($r=0.237$, $p>0.001$). **Conclusion:** Findings of the current study show that BMI is significantly correlated with severity of pain of knee in OA patients as higher the BMI greater the pain of knee. Therefore, pain in knee patients with higher BMI should be advised to decrease weight via suitable exercise program.

Keywords: Osteoarthritis (OA), Obesity, Numerical pain rating scale(NPRS).

INTRODUCTION

Osteoarthritis (OA) a well known degenerative disorder is one of the commonest forms of arthritis.^[1] Knee OA is a heterogeneous disorder having different etiological factors in different patients. Among most common causative factors are advancing age, female gender, alignment defects, genetic factors and obesity etc.^[3] Prevalence of knee pain has been found up to 50% in individuals above 50 years. Moreover, half of these patients suffer from severe and disabling knee pain.^[2] Obesity is emerging as an epidemic throughout the world. Obesity is the most important risk factors in the development of OA.^[2,3]

Obesity is a modifiable risk factor which can decrease the risk as well as the severity of knee pain in OA patients.^[4] Therefore, present study was designed to evaluate the correlation higher body mass index (BMI) with knee pain of OA patients.

MATERIALS AND METHODS

This was an observational type of study conducted in Department of Orthopaedic RMRI, Bareilly (U.P.). Total one hundred five patient of

OA with knee pain were included in this study. Inclusion criteria for the study were age ≥ 35 years and radiological evidence of knee OA with knee pain. Exclusion criteria for the study were: history of knee injury, smoking, hormonal therapy (corticosteroids, hyaluronidase, prolotherapy injections), HLA B 27, Gout. OA of knee was graded according to Kellgren and Lawrence grading.⁷ Simple linear regression analysis and Pearson correlation coefficient were used to evaluate if there is any association between BMI and numerical pain rating scale (NPRS). A p value <0.05 was considered statistically significant.

RESULTS

Total of one hundred and five patients (age 54.39 ± 13.17 years) were included in the present study. Among them 61 (58.08%) male patients and 44 (41.92%) were female patients. The subjects were comparable with respect of age, weight and BMI. Results are present as mean \pm SD.

[Table 1] show that different variables of the OA patients. The age, Weight and BMI of the OA patients were 50.58 ± 10.87 years, 66.84 ± 11.08 Kg and 1.57 ± 0.12 kg/m² correspondingly.

Name & Address of Corresponding Author

Dr. Atul Panday
Assistant Professor,
Department of Orthopaedics,
RMRI, Bareilly,
Uttar Pradesh, India.

Table 1: Demographic Parameters.

No. of patients (n=105)	Mean \pm SD
Age (yrs)	50.58 \pm 10.87
Weight (Kg)	66.84 \pm 11.08
Height (m)	1.57 \pm 0.12

[Table 2] reveals that among all OA patients 52% bilateral, 27.6% right and 20.4% left knee pain suffering. Further, it is evident from table 2 that number of OA patients 12 (11.4%), 40(38.09%), 30 (28.57%) and 33 (31.4%) were suffering with grade 1 to 4 OA severities respectively according to Kellgren and Lawrence grading. It is evident from table 2 that NPRS of the study ranged from 2 to 9 (mean 5.62±1.89).

[Table 3]. shows that there was a significant positive correlating of the NPRS score with BMI ($r=0.237$, $p<0.0001$). Further, regression analysis shows that NPRS can significantly estimated from respective BMI ($R^2 = 0.561$).

Table 2: Comparison of BMI and NPRS.

Grade	No.	BMI (Mean±SD)	NPRS (Mean±SD)
1	12	24.1±1.32	3.57±1.20
2	40	25.8.00±2.02	5.04±1.36
3	30	27.56±3.43	6.45±1.32
4	33	28.66±4.24	6.8±1.22
Total	105	26.89±3.94	5.62±1.89
Pearson correlation between BMI and NPRS	r	R²	P value
	0.237	0.561	>0.001

DISCUSSION

OA is one of the commonest degenerative disorders prevalent in elderly population. The prognosis and development of OA depends on various factors including structural, environmental and anatomical. Moreover, some additional factors influence the outcome of OA via dietary intake, level of physical activity and injury to the knee.

Findings of the present study revealed that there was a significant correlation of BMI with the severity of pain in OA patients. The BMI seems to be the most easily modifiable factor found in OA with knee pain. Previous research suggests that knee replacement surgeries include more than 50% overweight conditions.^[8] Berebaum F et al recorded that excessive vertical loading forces Over knee joint of patients suffering with OA can leads to inflammatory process.^[9] Further, Leptins have been found associated with chronic inflammatory state of knee joint in OA. Moreover, leptons found in abandoned quantity in adipose tissue.^[10] Therefore, female gender, obesity, tendency to body fat ratio may increase the severity of knee pain with OA.

Single leg stance exert up to six time body weight through knee joint, Further any increase in body weight may increase the transmitted force from knee joints into multiples. Moreover it can be evident from the knee of an overweight individual during walking. Increased BMI leads to progressive destruction of knee joints and induces pain which in turn results in decrease physical activity. In this way a dilemma developed in condition of as patients are unable to move due to pain which in

turn leads to further increase of body weight and worsen the condition of knee in OA patients.^[4-7]

Furthermore, prior study suggest that individuals with OA with radiological evidences of knee worsening have a severe pain in higher BMI in comparison of lower BMI patients. However, findings of that study also suggested that individual with higher BMI experience greater pain in knee compare to lower BMI.^[5] Findings of the current study recorded that BMI alone may account for 56.1% of total variations of NPRS severity. This finding is consistent with the finding of the previous study of Matthew et al as they observed that pain subjects showed a higher BMI (30.4 kg/m²) in comparison of without pain individuals (27.5 kg/m²) ($p < 0.0001$). Moreover, Severity of pain has been found increased with every successive BMI group. This study concluded that BMI is positively correlated with NPRS in OA patients with knee pain.

CONCLUSION

Findings of the current study show that BMI is significantly correlated with severity of pain of knee in OA patients as higher the BMI greater the pain of knee. However, decreasing the BMI may reverse the OA changes and decrease the severity of pain. Therefore, OA patients with knee and higher BMI should be advised to reduce weight via suitable exercise program. One can consider about bariatric treatment options if BMI is too high for early and quick results.

REFERENCES

1. Riis A, Rathleff MS, Jensen MB, et al. Low grading of the severity of knee osteoarthritis pre-operatively is associated with a lower functional level after total knee replacement: a prospective cohort study with 12 months' follow-up. *Bone Joint J* 2014; 96-B:1498-502.
2. Yoon KH, Lee JH, Kim JW, et al. Epidemic obesity and type 2 diabetes in Asia. *Lancet* 2006;368:1681-8.
3. Felson DT, Zhang Y, Hannan MT, Naimark A, Weissman B, Aliabadi P, Levy D: Risk Factors for Incident Radiographic Knee Osteoarthritis in the elderly: the Framingham Study. *Arthritis Rheum* 1997, 40:728-33
4. Lee J, Song J, Hootman JM et al. Obesity and other modifiable factors for physical inactivity measured by accelerometer in adults with knee osteoarthritis: data from the Osteoarthritis Initiative (OAI). *Arthritis Care Res* 2013; 65:53-61.
5. Weiss E. Knee osteoarthritis, body mass index and pain: data from the Osteoarthritis Initiative; *Rheumatology* 2014; 53:2095-2099.
6. Rogers M W, Wilder F V. The association of BMI and knee pain among persons with radiographic knee osteoarthritis: A cross-sectional study; *BMC Musculoskeletal Disorders* 2008, 9:163.
7. Kellgren, JH.; Lawrence, JS. Atlas of Standard Radiographs of Arthritis. II. Philadelphia, PA: F.A. Davis; 1963. The epidemiology of chronic rheumatism.
8. D'Arcy Y . Pain and obesity. *Nurs Manage* 2012;43:21-6

9. Berenbaum F, Eymard F, Houard X. Osteoarthritis, inflammation and obesity. *Curr Opin Rheumatol* 2013;25: 114-8
10. Gaumont I, Arsenault P, Marchand S. The role of sex hormones on formalin-induced nociceptive responses. *Brain Res* 2002;958:13

How to cite this article: Saxena DK, Panday A. Body Mass Index and its Correlation with Knee Pain in Osteoarthritis Patients. *Ann. Int. Med. Den. Res.* 2018; 4(4):OR15-OR17.

Source of Support: Nil, **Conflict of Interest:** None declared