

# Comparative Evaluation of Nemoceph and Foxit PDF Reader for Steiner's Cephalometric Analysis.

Mukesh Kumar<sup>1</sup>, Pramod Shetty<sup>2</sup>, Prakrathi Shetty<sup>3</sup>, Sommya Kumari<sup>4</sup>, Rameshwar Singh<sup>5</sup>

<sup>1</sup>Assistant Professor, Faculty of Dentistry, Vardhman Institute of Medical Sciences, Pawapuri, Nalanda, Bihar.

<sup>2</sup>Professor, Pacific Dental College, Udaipur, Rajasthan.

<sup>3</sup>B.D.S.

<sup>4</sup>Post Graduate Student, Post Graduate Student, Buddha Institute of Dental Sciences and Hospital, Patna.

<sup>5</sup>Prof & Head, Vardhman Institute of Medical Sciences, Pawapuri.

Received: March 2019

Accepted: April 2019

**Copyright:** © the author(s), publisher. Annals of International Medical and Dental Research (AIMDR) is an Official Publication of "Society for Health Care & Research Development". It is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

## ABSTRACT

**Background:** Cephalometry is used to quantify and qualify the type of malocclusion. The digital technology has overcome the limitations of manual method. A high cost of the commercially available software is unaffordable for its wide spread use. **Objectives:** The aim of this study was to evaluate and compare the values of Steiner's cephalometric analysis using Nemoceph and Foxit PDF reader. No significant difference between the two methods will result in that Foxit PDF reader can be used as an cost effective alternative. **Methods:** This study was conducted on 100 digital lateral cephalogram taken from the same machine. The samples were collected by non-probability convenience sampling procedures. These images were analyzed for Steiner's Cephalometric Analysis using the two software. **Results:** The skeletal and dental values showed no statistical significant difference in the majority, except for the L1-NA (Linear) and L1-NB (Linear). **Conclusion:** Results showed that there is a high agreement between the two methods.

**Keywords:** Cephalometry; Digital Imaging ; Software.

1

## INTRODUCTION

Harmonious facial esthetics and optimal functional occlusion with a firm structural balance is a recognized canon for orthodontists.<sup>[1,2]</sup> A scientific approach to analyze the human craniofacial patterns was pioneered by anthropologists and anatomists.<sup>[3-5]</sup> Since the introduction of cephalometric radiography by Broadbent in 1931, significant advancement has been achieved over the years.<sup>[4-7]</sup> The vital role of cephalometric analysis in orthodontic diagnosis, treatment planning, and monitoring treatment and growth changes is well established.<sup>[7-9]</sup>

The traditional hand-tracing process of cephalometric analysis uses an acetate overlays, pencil, ruler and protractor to measure the linear and angular values. Though most economical and accessible, the potential systematic and random error (Bregersen 1980, Houston 1983), high time demand, special dark chamber, chemical hazard, together with difficult archiving are amongst the possible cause of its set back.<sup>[10-11]</sup>

Digital radiographic technique emerged During the 1980s and early 1990s. These digital cephalometric

images created a surge for computer cephalometric software.<sup>[12]</sup> This technological advancement not only overcome the limitations of the conventional technique but enabled brightness and contrast control facilitating easy landmark identification for accuracy.<sup>[13-16]</sup>

The availability, and affordability and user friendly score of these commercially available software remained questionable. (Nouri M. et.al. 2015).<sup>[17]</sup> Therefore, the present study was conducted with an objective to compare and evaluate the measurement obtained using NEMOCEPH (a commercially available software for cephalometric analysis) and the measurement tool of FOXIT PDF READER for STEINER'S Cephalometric Analysis.

## MATERIALS & METHODS

Soft copy images of digital lateral cephalograms of 100 prospective orthodontic patients were include in the study. All the radiographs were taken from the same digital OPG machine with an automatic KVp and mA setting. The samples were selected through a non-probability convenience sampling procedures. All the radiographs were selected on the basis of quality and clarity of images and with ease for identification of landmarks. The selection criteria were not be affected by age, gender, machine, head positioning and tooth contact. Poor quality image, distortion, artifact and craniofacial anomalies were excluded from the study.

### Name & Address of Corresponding Author

Prof. (Dr.) Pramod Shetty  
Professor,  
Pacific Dental College,  
Udaipur, Rajasthan.

The lateral cephalometric images were cropped to the size of standard lateral cephalogram film (8x10 inch) using Adobe Photoshop. A ruler scale image of 8 inch was added on the top of this image, extending from the right margin to the left margin for easy calibration with the software to be tested. The standardized and calibrated images were numbered 1-100 on the upper right hand side corner of the images for identification. The images were saved in JPEG format, with maximum quality setting at 200 dpi.

The Landmark identification was carried using - NEMOCEPH NX 2009 for Windows and FOXIT PDF READER, version 3.0 (a free download software)

A laptop with a mouse-controlled cursor was used for on screen landmark identification and cephalometric analysis using NemoCeph NX 2009 for Windows (Nemotec, Madrid, Spain). The cephalometric image to be evaluated was marked for the cephalometric landmarks as demanded by the software. The same laptop with a mouse-controlled cursor was used for on screen cephalometric analysis using the measurement tools in the tool bar of

FOXIT PDF READER software. The “Distance Tool” and the “Area Tool” were used for the linear and the angular measurements and immediately the observed values were recorded manually on a paper. Image magnification and contrast enhancement tool was used for easy identification of landmarks.

The data was subjected to statistical analyses using Statistical Package for Social Sciences Software version 11.0 (SPSS Inc, Chicago, IL). Oneway ANOVA was used for comparison between the variables and Post Hoc test followed by Turkey’s test was done to check the level of significance.

## RESULTS

The pre-standardized and pre-calibrated digital lateral cephalometric radiographs evaluated for the five skeletal and five dental values of Steiner’s Analysis using the FOXIT PDF READER and NEMOCEPH showed – The mean difference of the skeletal values obtained using the two software showed no significant statistical difference [Table 1 & 2].

**Table 1: Comparison Of Skeletal Values Between Groups (One way ANOVA)**

Variable	Group	Mean value	S.D	p-value
SNA	Nemoceph	82.83	4.45	0.79
	Foxit	82.16	4.97	
SNB	Nemoceph	78.00	5.48	0.95
	Foxit	77.83	6.09	
ANB	Nemoceph	5.15	3.15	0.69
	Foxit	5.53	3.93	
Mandibular Plane Angle	Nemoceph	28.95	7.69	0.80
	Foxit	27.73	9.03	
Occlusal Plane Angle	Nemoceph	15.85	6.31	0.51
	Foxit	14.28	5.49	

(\*p ≤ 0.5 Significant).

**Table 2: Comparison Of Skeletal Values Between Groups (Post Hoc Test)**

Variable	Group	Group	Mean difference	p-value
SNA	Nemoceph	Foxit	0.67	0.80
SNB	Nemoceph	Foxit	0.17	0.99
ANB	Nemoceph	Foxit	0.38	0.87
Mandibular Plane Angle	Nemoceph	Foxit	1.23	0.78
Occlusal Plane Angle	Nemoceph	Foxit	1.57	0.49

(\*p ≤ 0.5 Significant).

**Table 3: Comparison of Dental values between Groups (One way ANOVA)**

Variable	Group	Mean value	S.D	p-value
U-1 to NA (Angle)	Nemoceph	24.98	9.44	0.34
	Foxit	28.23	10.79	
U-1 to NA (Linear)	Nemoceph	0.22	0.15	0.0001
	Foxit	6.55	4.21	
L-1 to NB (Angle)	Nemoceph	29.83	6.87	0.12
	Foxit	29.85	8.33	
L-1 to NB (Linear)	Nemoceph	0.28	0.11	0.0001
	Foxit	7.29	2.85	
Inter Incisal Angle	Nemoceph	120.29	11.93	0.73
	Foxit	118.59	10.92	

(\*p ≤ 0.5 Significant).

**Table 4: Comparison of Dental values between Groups (Post Hoc Test)**

Variable	Group	Group	Mean difference	p-value
U-1 to NA (Angle)	Nemoceph	Foxit	3.25	0.33
U-1 to NA (Linear)	Nemoceph	Foxit	6.33	0.0001
L-1 to NB (Angle)	Nemoceph	Foxit	0.02	1.00
L-1 to NB (Linear)	Nemoceph	Foxit	7.02	0.0001
Inter Incisal Angle	Nemoceph	Foxit	1.70	0.83

(\*p ≤ 0.5 Significant).

The dental values showed no statistical significant difference in the majority, except for the L1-NA (Linear) and L1-NB (Linear). (\* $p \leq 0.5$  Significant). [Table 3 and 4]

## DISCUSSION

A precise diagnosis and treatment planning is essential to the success of orthodontic treatment. In 1931, Orthodontics ushered in the age of radiographic cephalometry.<sup>[18]</sup> Since then, the orthodontic domain has achieved a new horizon both in research and clinical science.<sup>[19,20]</sup> A number of different cephalometric analyses and norms are available today.<sup>[21,22]</sup>

Traditional cephalometric radiography and analysis were done manually using a large inventory and was prone to errors. The technique also is laden with weakness.<sup>[23,24]</sup>

With the rapid evolution of digital radiography landmark location and on screen tracing has become area of interest for researchers. Computer-aided cephalometric analysis on digitized cephalogram substantially reduces the potential errors, eliminate the production of hard copies and is time saving as well. Currently, cephalometric analyses for orthodontic diagnosis, treatment planning, and research are often performed on digital images by means of computer software.<sup>[25,26]</sup> The high cost and availability account limitation of these software programs.

The innovative techniques of Prawat J.S. et. al. (1995), Shahidi S. et.al.; (2013) and Nouri M. et.al.; (2015) have attempted cost effective options with success.<sup>[17,27,28]</sup>

Precision and reproducibility in data is an essential requirement.<sup>[7]</sup> Duraó et.al.; (2015) reported a lower level of reproducibility in landmarks identification among orthodontists compared to maxillofacial radiologist.<sup>[29]</sup>

The current study compared the NEMOCEPH and the measurement tools in the tool bar of FOXIT PDF READER for cephalometric analysis. The pre-standardized and pre-calibrated digital lateral cephalometric radiographs evaluated for the five skeletal and five dental values of Steiner's Analysis revealed – the mean difference of the skeletal values obtained using the two software showed no significant statistical difference. This was in consonance with the study reports of Erkan M., et. al.; (2012), Goracci C. and Ferrari M. (2014), Rusa O et. al. (2015) and CORREIA T.R.G.S. (2017) who reported a high consistency between the different software evaluated.<sup>[30-32]</sup>

T Sommer et. al. (2009) reported a difference of below 2° but clinically acceptable for the mid face structures. The dental values in our study showed no statistical significant difference in the majority, except for the L1-NA (Linear) and L1-NB (Linear). A similar finding was reported by Celik E. et. al.; (2009) and ALDREES A. M.(2010). Study report

of Tsorovas G. and Karsten A.L.(2010) showed better agreement with the advanced features of software for [(L1 to NB (mm)], compared to standard feature.<sup>[33-36]</sup>

The overall finding of our study showed 80 percent agreement between the two methods. Therefore, FOXIT PDF READER can be used as an alternative to commercially available NEMOCEPH software.

## CONCLUSION

Orthodontics is undergoing a gradual transition and has reached a digital era. The specialty has experienced advancements both in techniques and technology. The present study has shown a comparable and non significant difference between the data obtained with the two softwares, except for a few exceptions. Therefore, FOXIT PDF READER can be considered as a cost effective alternative.

## REFERENCES

1. Kommi PB, Venkatesan R, Keerthi N, Kumar AN, Kumar S, Gopinath V. A Cephalometric Assessment of Ideal Nasolabial Angle Range for South Indian Population. *Journal of International Oral Health*. 2016; 8(2): p. 205-207.
2. Soni A, Alladwar N, Goel S, Chopra R, Sharma S. Evaluation of lateral Cephalometric Norms for Burstone's Analysis in Chhattisgarh by using Nemoceph Software with Lateral Cephalograms Taken in Natural Head Position. *International Journal of Oral Health Dentistry*. 2015; 1(3): p. 114-119.
3. Turkdonmez CO, Taner L, Uzuner FD, Gungor K. Craniofacial Evaluation of Class I Turkish Adults: Bimler Analysis. *Turkish Journal of Orthodontics*. 2014; 26(4): p. 169-176.
4. Turkdonmez CO, Taner L, Uzuner FD, Gungor K. Craniofacial Evaluation of Class I Turkish Adults: Bimler Analysis. *Turkish Journal of Orthodontics*. 2014; 26(4): p. 169-176.
5. Gribel BF, Gribel MN, Frazao DC, McNamara Jr JA, Manzi FR. Accuracy And Reliability Of Craniometric Measurements On Lateral Cephalometry And 3D Measurements On CBCT Scans. *The Angle Orthodontist*. 2011; 81: p. 28-37.
6. Naragond A, Kenganal S, Sagarkar R, Kumar S, Day S. Diagnostic Limitations of Cephalometrics in Orthodontics-A Review. *Journal of Dental and Medical Sciences*. 2012; 3(1): p. 30-35.
7. Ganna PS, Shetty SK, Yethadka MK, Ansari A. An Evaluation of the Errors in Cephalometric Measurements on Scanned Lateral Cephalometric Images using Computerized Cephalometric Program and Conventional Tracings. *Journal Of Indian Orthodontic Society*. 2014; 48(4): p. 388-392.
8. Anh TT, Dang TV, Hong An NP, Ngoc VTN, Phuong NTT, Anh LQ. Cephalometric Norms For The Vietnamese Population. *APOS Trends in Orthodontics*. 2016; 6(4): p. 200-204.
9. Puralmal K, Alam MK, Zam Zam NM. Cephalometric Norms of Malaysian Adult Chinese. *International Medical Journal*. 2013; 20(1): p. 87-91.
10. Da Silva JMG, De Melo Castilho JC, Matsui RH, Matsui MY, Gomes MF. Comparative Study Between Conventional And Digital Radiography In Cephalometric Analysis. *Journal of Health Science Institution*. 2011; 29(1): p. 19-22.
11. Gupta G, Tripathi AA, Khandekar S. Cephalometric VTO : A Blueprint. *Heal Talk*. 2013; 05(3): p. 47-50.

12. Rodrigues CD, Da Silveira MMF, Tavano O, Shibuya RH, Modesto G, Estrela C. Evaluation Of Indirect Methods Of Digitization Of Cephalometric Radiographs In Comparison With The direct Digital Method. Dental Press Journal of Orthodontics. 2010; 15(4): p. 124-132.
13. Oshagh M, Shahidi S, Momeni DS. Effects Of Image Enhancement On Reliability Of Landmark Identification In Digital Cephalometry. Indian Journal of Dental Research. 2013; 24(1): p. 98-103.
14. Segura FJE, Valverde AS, Ocampo AM, Angelares PRC. Comparative Study Between Digital And Manual Cephalometry With Digital Radiographs. Revista Mexicana de Ortodoncia. 2014; 2(2): p. 93-96.
15. Sayinsu K, Isik F, Trakyali G, Arun T. An evaluation of the errors in cephalometric measurements on scanned cephalometric images and conventional tracings. European Journal of Orthodontics. 2007.; 29: p. 105-108.
16. Grybauskas S, Balciuniene I, Vetra J. Validity and Reproducibility of Cephalometric Measurements Obtained from Digital Photographs of Analogue Headfilms. Stomatologija Baltic Dental and Maxillofacial Journal. 2007; 9(4): p. 114-120.
17. Nouri M, Hamidiaval S, Baghban AA, Basafa M, Fahim M. Efficacy of a Newly Designed Cephalometric Analysis Software for McNamara Analysis in Comparison with Dolphin Software. Journal of Dentistry. 2015; 12(1): p. 60-69.
18. Paixao MB, Sobral MC, Vogel Cj, De Araujo TM. Comparative Study Between Manual And Digital Cephalometric Tracing Using Dolphin Imaging Software With Lateral Radiographs. Dental Press Journal of orthodontics. 2010; 15(6): p. 123-130.
19. Erkan M, Gurel HG, Nur M, Demirel B. Reliability of Four Different Computerized Cephalometric Analysis Programs. European Journal of Orthodontics. 2012; 34(3): p. 318–321.
20. Kumar V, Sundareswaran s. Cephalo metric assessment of sagittal dysplasia: a review of twenty one methods. The Journal Of Indian Orthodontic Society. 2015; 48(1): p. 33-41.
21. Cutoviu T, Joviu N, Kozomara R, Radojipiu J, Janoseviu M, Mladenoviu I, et al. Cephalometric analysis of the middle part of the face in patients with mandibular prognathism. Vojnosanitetski Pregled. 2014; 71(11): p. 1026–1033.
22. Purnal K, Alam MK, Zam Zam NM. Cephalometric Norms of Malaysian Adult Chinese. International Medical Journal. 2013; 20(1): p. 87-91.
23. McNamara JA. A Method of Cephalometric Evaluation. American Journal of Orthodontics. 1984; 86(6): p. 449-469.
24. Naumova J, Lindman R. A Comparison of Manual Traced Images and Corresponding Scanned Radiographs Digitally Traced. European Journal of Orthodontics. 2009; 31(3): p. 247– 253.
25. Kumar D, Solanki R, Sharma R, Khangwal M, Berwal V, Gupta S. Cephalometric Landmark Identification by Conventional and Digital Direct Radiography in Skeletal Class II Individuals. International Journal of All Research Education and Scientific Methods. 2015; 3(12): p. 6-10.
26. Oshagh M, Shahidi S, Momeni Danaei S. Effects Of Image Enhancement On Reliability Of Landmark Identification In Digital Cephalometry. Indian Journal of Dental Research. 2013; 24(1): p. 98-103.
27. Prawat JS, Nieberg L, Cisneros GJ, Acs. G. A Comparision Between Radiographically And Sonically Produced Cephalometric Values. Angle orthodontics. 1995; 65(4): p. 271-276.
28. Shahidi S, Oshagh M, Gozin F, Salehi P, Danaei S. Accuracy of Computerized Automatic Identification of Cephalometric Landmarks by a Designed Software. Dentomaxillofacial Radiology. 2013; 42: p. 1-8.
29. Durao APR, Morosolli A, Pittayapat P, Bolstad N, Ferreira AP, Jacobs R. Cephalometric landmark variability among orthodontists and dentomaxillofacial radiologists: a comparative study. Imaging Science in Dentistry. 2015; 45: p. 213-220.
30. Goracci C, Ferrari M. Reproducibility of measurements in Tablet-assisted, PC-Aided, and Manual Cephalometric Analysis. Angle Orthodontics.. 2014; 84: p. 437–442.
31. Rusu OL, Petcu AE, Dragan E, Haba D, Moscalu M, Zetu IN. Reliability and Accuracy of Three Different Computerized Cephalometric Analysis Software. Revista Medico-Chirurgicala a Societatii De Medici Si Natralisti Din Iasi. 2015; 119(1): p. 248-256.
32. Correia ,TRGS, Sena LMFDSJSPD, Pereira HSG. Cephalometric Analysis: Concordance Between Software.. Rev Gauch Odontol, Porto Alegre. 2017; 65(4): p. 321-325..
33. Celik E, Polat-Ozsoy O, Memikoglu TUT. Comparison of Cephalometric Measurements with Digital Versus Conventional Cephalometric Analysis. European Journal of Orthodontics. 2009; 31: p. 241– 246.
34. Sommer T, Ciesielski R, Erbersdobler J, Orthuber W, Fischer-Brandies H. Precision of Cephalometric Analysis via Fully and Semiautomatic Evaluation of Digital Lateral Cephalographs. Dentomaxillofacial Radiology. 2009; 38: p. 401–406.
35. Tsorovas G, Karsten ALA. A Comparison of Hand-Tracing and Cephalometric Analysis Computer Programs With and Without Advanced Features — Accuracy and Time Demands.. European Journal of Orthodontics. 2010; 32: p. 721–728..
36. ALDREES AM. Evaluation of Incisors Inclination in Five Cephalometric Analysis Methods. Pakistan Oral & Dental Journal. 2010; 30(2): p. 1-8.

**How to cite this article:** Kumar M, Shetty P, Shetty P, Kumari S, Singh R. Comparative Evaluation of Nemoceph and Foxit PDF Reader for Steiner's Cephalometric Analysis. Ann. Int. Med. Den. Res. 2019; 5(3):DE64-DE67.

**Source of Support:** Nil, **Conflict of Interest:** None declared