

# Comparison of Two Different Doses of Dexmedetomidine as an Adjuvant to Intrathecal Bupivacaine in Patients Undergoing Femur Surgery.

Gunjan Chaudhry<sup>1</sup>, Pranav Bansal<sup>2</sup>, Deepak Singla<sup>1</sup>, Jagdish Dureja<sup>3</sup>, Surya Dahiya<sup>4</sup>

<sup>1</sup>Assistant Professor, Dept. of Anaesthesiology, BPS Govt. Medical College for Women, Khanpur, Sonapat, Haryana.

<sup>2</sup>Associate Professor, Dept. of Anaesthesiology, BPS Govt. Medical College for Women, Khanpur, Sonapat, Haryana.

<sup>3</sup>Professor and Head, Dept. of Anaesthesiology, BPS Govt. Medical College for Women, Khanpur, Sonapat, Haryana.

<sup>4</sup>Senior Resident, Dept. of Dentistry, BPS Govt. Medical College for Women, Khanpur, Sonapat, Haryana.

## ABSTRACT

**Background:** Duration of action of local anaesthetic is an important limiting factor in spinal anaesthesia. Dexmedetomidine, selective  $\alpha$  2- agonist has been recently used in addition to other adjuvants to prolong the duration of intrathecal local anaesthetics. **Aim:** To compare two different doses of dexmedetomidine added to heavy bupivacaine 0.5% intrathecally for femur surgeries. **Methods:** In this prospective double blind trial, one hundred and twenty patients were randomly allocated into two groups, D1 and D2. Group D1 received 12.5 mg 0.5% hyperbaric bupivacaine and 5  $\mu$ g dexmedetomidine. Group D2 received 12.5 mg 0.5% hyperbaric bupivacaine and 10  $\mu$ g dexmedetomidine. **Results:** Sensory and motor block were comparable in both groups in terms of characteristics like the highest level of sensory block achieved, time to achieve maximum sensory block and time of two segment regression, time to achieve maximal motor block and duration of regression to Bromage scale 0. However time of first analgesic request and total analgesic requirement were significantly reduced by increasing intrathecal dose of dexmedetomidine to 10  $\mu$ g without any undesirable effects. **Conclusion:** Intrathecal dose of 10  $\mu$ g dexmedetomidine provided an increased duration of sensory compared to 5 $\mu$ g dosing, with no significant increase in duration of motor blockade or the incidence of hypotension, bradycardia and any other undesirable side-effects.

**Keywords:** Analgesic,  $\alpha$ -2 agonist, Intrathecal Dose, Motor Blockade, Spinal Anaesthesia.

## INTRODUCTION

Orthopaedic fixations in lower extremity has conventionally and preferable been performed under spinal anaesthesia. Though, cost effective and easy to perform, the major limitation observed frequently is the limited duration of action, which requires conversion of anaesthetic technique or early analgesic intervention in post-operative period.<sup>[1]</sup> The list of various intrathecal adjuvants that have been used for prolongation and intensifying the effects of spinal anaesthesia includes opioids like fentanyl, morphine, clonidine, neostigmine, and more recently dexmedetomidine.<sup>[2-4]</sup>

However, most of these adjuvant drugs are associated with their own set of side effects, which limits their use. For example, opioid additives are associated with urinary retention, pruritus or delayed respiratory depression (more frequently with buprenorphine) while neostigmine is associated with a high incidence of nausea and vomiting.<sup>[5]</sup>

and analgesia. Data on its utility in central neuraxial blockade is limited with only a few studies on extensive medline and google search. Dexmedetomidine provides faster onset of both sensory and motor blockage, increased duration of sensory and motor blockade, with no significant side effects at smaller clinical doses being used.<sup>[4,6]</sup> A few human studies have earlier shown that addition of intrathecal dexmedetomidine to hyperbaric bupivacaine in spinal anaesthesia increased the duration of analgesia without significant side effects. Yet, the ideal intrathecal dose of dexmedetomidine is still a matter of debate.<sup>[4,6,7]</sup>

It is well known that surgeries on femur bone are not only painful but sometimes are prolonged in duration that they run short of peri-operative or immediate post-operative analgesia. In the current study, we aimed to compare two different doses of dexmedetomidine added to heavy bupivacaine 0.5% intrathecally for fracture femur surgeries

## MATERIALS AND METHODS

In this prospective, double blind trial, after approval from institutional review board, 120 patients of ASA I & II status, between age 18-60 years and both genders, scheduled for fixation on fracture femur, during the study period from Jan 2014 to March 2015 (15 months) were enrolled for study. Written informed consent was obtained from all patients. Patients with any contraindication to spinal anaesthesia, hemodynamic instability,

### Name & Address of Corresponding Author

Dr Gunjan Chaudhry  
Assistant Professor,  
Dept. of Anaesthesiology,  
BPS Govt. Medical College for Women, Khanpur,  
Sonapat, Haryana, India.  
E mail: gunjan26117@gmail.com

Dexmedetomidine, a new selective  $\alpha$  2-agonist, is a non-opioid which has been used intravenously for providing peri-operative supplementary sedation

## ***Chaudhry; Comparison of Different Doses of Dexmedetomidine as an Adjuvant to Intrathecal Bupivacaine in Femur Surgery***

history of coagulopathy or associated head injury with GCS scores < 15 were excluded from the study.

Patients were evaluated in the pre- anaesthetic clinic prior to surgery and were explained about the procedure of spinal anaesthesia, visual analogue scale of pain and for their willingness to participate in the study.

On the day of surgery, the baseline VAS score was recorded in a pre - operative room in all the cases. Patients were randomized into two groups D1 and D2 using computer generated sequences. A two-operator technique was employed to maintain blinding where the operator preparing the drugs according to the group allocation was not involved in further patient management. In the operating room, standard monitoring that included pulse rate, SpO<sub>2</sub>, non-invasive blood pressure and ECG was attached and lactated Ringer's solution of 500 ml was infused over 15 to 20 mins. With the patient in the sitting position a lumbar puncture was performed at L3-L4 inter space using midline approach by a 26-gauge Quincke needle. Cases in group D1 received 12.5 mg 0.5% hyperbaric bupivacaine and 5 µg dexmedetomidine while in group D2 received 12.5 mg 0.5% hyperbaric bupivacaine and 10 µg dexmedetomidine. Further interventions and monitoring were done by an investigator blinded to the group allocation. After spinal anaesthesia, all patients were placed in supine position and a level of T10 dermatome was achieved. Sensory level was assessed using loss of pinprick sensation in midline while modified Bromage scale was used to assess motor block. Time taken to achieve the desired effect was noted prior to surgery.

After surgery VAS scale was recorded before shifting the patient out of the operation theatre, at half hour intervals during 1<sup>st</sup> hour and thereafter at every hourly for next 8 hours and at 12 and 24 hours. Duration of two sensory level regression and

regression of motor power to Bromage scale reading of 0 were also recorded. Duration of pain relief was measured from time of spinal anaesthetic injection to first request of rescue analgesics. Rescue analgesics used were inj. Diclofenac 75 mg BD. If satisfactory analgesia was not achieved than inj. Paracetamol 1 gm QID was added.

The incidence of side effects like nausea, vomiting, bradycardia and hypotension were recorded in both the groups and managed as per the standard guidelines.

### **Statistical Analysis**

To detect a 20% difference in the response rate among the groups with a minimum response of 50% estimated from initial pilot observations, with 90% power and 5% alpha error a sample size of 60 cases per group was required. The sample size was calculated using the power and sample size calculator of the Department of Biostatistics, Vanderbilt University, US.

Analysis of data was done using SPSS® version 16 (Statistical Packages for the Social Sciences, Chicago, IL). Results are expressed as mean and standard deviation (SD), median and range or numbers. Quantitative data were compared using one-way analysis of variance (ANOVA), and independent samples *t*-test and qualitative data was analysed using chi square test. *The P value* of <0.05 was considered statistically significant.

## **RESULTS**

A total of 120 ASA grade I, II patients were studied and the data collected was analysed. Patients were divided into two groups with 60 patients each. Both groups were comparable in terms of age, height, weight, ASA status, and duration of surgery as shown in [Table 1].

**Table 1:** Patient Characteristics.

	<b>D1 (Mean ± S.D.)</b>	<b>D2 (Mean ± S.D.)</b>	<b>P-Value</b>
Number of cases	60	60	--
Age (yrs)	34.57 (± 5.33)	34.68 (± 4.88)	0.073
Height (cms)	159.53 (± 4.54)	157.89 (± 5.03)	0.065
Weight (Kgs)	62.30 (± 3.64)	61.95 (± 2.75)	0.084
ASA Status ASA I	45	43	0.072
ASA Status ASA II	15	17	0.069
Duration of surgery	79.7 (54-126)	82.05(47- 138)	0.059

Similarly other variables like baseline haemodynamic data and pain scores didn't differ significantly in both the groups.

Characteristics of sensory block were compared and no significant difference was noted in terms of

highest level of sensory block achieved, time to achieve maximum sensory block and time of two-segment regression expressed in minutes [Table 2].

**Table 2:** Characteristics of Sensory Block.

Block Characteristics	D 1(Mean ± S.D.)	D 2(Mean ± S.D.)	P-Value
Highest level of sensory block	T6-T10	T6-T8	NS
Time (min) to achieve sensory block	16.40 ± 4.45	15.40 ± 4.30	0.724
Time to achieve two-segment regression (min)	107.30 ± 20.05	115.35 ± 25.05	0.051
Time to achieve maximum Bromage scale motor block (min)	18.10 ± 4.10	19.05 ± 4.35	0.065
Duration of regression of motor block to modified bromage 0	221.40 ± 35.60	231.40 ± 45.50	0.053
Time to first analgesic request (min)	210.30 ± 45.20	254.18 ± 40.55	0.029
Number of doses of diclofenac in the first 24 hours	84	68	0.037
Hypotension	3	5	0.071
Bradycardia	5	5	--

Similarly, motor block characteristics, i.e. time to achieve maximal motor block and duration of regression to Bromage 0 showed no significant difference [Table 2].

However, when groups were compared in terms of time for requirement of first analgesic and total requirement of analgesic significant differences were observed. Time from lumbar puncture to request for first analgesic was 210 ± 45.20 minutes for the D1 group while it was 245 ± 40.55 for D2 group. Similarly, total no. of analgesic dose requirement is significantly less (68) in the D2 group as compared to D1 (84) [Table 2]. This proved that increasing the dose of dexmedetomidine increases the duration of analgesia and has a post operative analgesic sparing effect.

Heart rate and blood pressure values were comparable in both groups. In group D1 3 cases and 5 cases in group D2 received ephedrine for control of hypotension. In each group, 5 patients required Inj atropine for control of bradycardia during the perioperative period. Nausea was reported in 2 patients in group D1 and 1 patient in group D2 but vomiting was not reported. Similarly, there was no incidence of respiratory depression, severe sedation or itching requiring active medical attention.

## DISCUSSION

Duration of action of intrathecally administered local anaesthetics is a major limiting factor for spinal anaesthesia. Various agents have been tried as adjuvant to prolong the effect of spinal anaesthesia. Dexmedetomidine, a dextro-isomer of medetomidine, is a highly selective  $\alpha$  2-drenoreceptor agonist. It has been shown that its affinity for  $\alpha$  2-adrenoreceptor is about ten times higher than that of clonidine.<sup>[8, 9]</sup> Activation of  $\alpha$  2-drenoreceptor has sympatholytic effect causing

bradycardia, hypotension, sedation and analgesia.<sup>[10]</sup>

Clonidine has already been shown to be an effective adjuvant to intrathecal local anaesthetics and opioids<sup>[10]</sup>. Comparative analysis of low dose dexmedetomidine (3  $\mu$ g) with clonidine (30  $\mu$ g) for intrathecal use has already been done by Kanazi et al<sup>[7]</sup> and it was found to be equipotent to a ten times dose of clonidine. However, the use of dexmedetomidine as intrathecal adjuvant has not been studied extensively.

In the above study, patients in the group that received 10  $\mu$ g dexmedetomidine had reduced postoperative pain scores and a longer analgesic duration than those who received 5  $\mu$ g dexmedetomidine as adjuvant to spinal bupivacaine. This shows that the use of dexmedetomidine as intrathecal adjuvant causes statistically significant prolongation of duration of spinal anaesthesia, decreased postoperative pain and lesser analgesic requirements. Also, the effect appears to be dose dependent with 10  $\mu$ g dexmedetomidine being more effective than 5  $\mu$ g with no significant increase in incidence of hypotension or bradycardia as well as any other undesirable side-effects.

Dexmedetomidine acts on  $\alpha$  2-adrenoreceptors located in the pre-synaptic C-fibres and post-synaptic dorsal horn neurons in the spinal cord. It acts on C-fiber inhibiting the release of pro-excitatory nociceptive transmitters i.e. glutamate & substance-P and hyper polarizes of post-synaptic dorsal horn neurons via G protein, mediated potassium channels resulting in analgesic effect.<sup>[11-12]</sup>

No significant increase in incidence of undesirable side effects like hypotension, bradycardia and sedation was noted in our study. Lack of effect on blood pressure can be due to the fact that local anaesthetics decrease sympathetic outflow themselves resulting in reduction of blood pressure.

So intrathecal dexmedetomidine cannot cause further hypotension probably because sympathetic inhibition maximally achieved by bupivacaine.<sup>[13]</sup> Lack of excessive sedation may be because CNS receptors responsible for this effect are spared from excessive doses of drug.<sup>[14]</sup> However sedation may occur due to systemic absorption or upward migration of drug when used in higher doses as seen in study by Al-Ghanem *et al*<sup>[4]</sup> who noted excessive sedation in 3 out of 25 patients. In our study, none of the patients reported excessive sedation or respiratory depression at 10 µg intrathecal dose of dexmedetomidine.

Animal studies have already been performed to assess safety of dexmedetomidine for intrathecal use and showed no adverse neurotoxic effects.<sup>[12,15,16]</sup> These studies also showed that epidural analgesic effect of α 2-adrenoreceptor agonist is directly related to their binding affinity.<sup>[17]</sup>

Though, our current knowledge regarding use of dexmedetomidine as intrathecal adjuvant has been improved by this study, it has certain obvious limitations. Firstly, only healthy individuals ASA I and II were studied, so the effect on older patients with cardiovascular comorbidities needs to be evaluated in future trials. Moreover, increasing doses of dexmedetomidine can be added intrathecally in future studies to observe the its desirable and undesirable effects.

## CONCLUSION

An Intrathecal dose of 10 µg dexmedetomidine can be safely used as an intrathecal adjuvant for increasing the duration of post-operative analgesia and decreasing post-operative analgesic requirements, compared to 5 µg dosing without any noticeable side effects.

## REFERENCES

1. Local anesthetics. In: Collins VJ, editor. Principles of Anesthesiology. 3rd ed, Vol. 2. Philadelphia: Lea and Febiger; 1993. p. 1232-81.
2. Boussofara M, Carlès M, Raucoules-Aimé M, Sellam MR, Horn JL. Effects of intrathecal midazolam on postoperative analgesia when added to a bupivacaine-clonidine mixture. *Reg Anesth Pain Med.* 2006;31:501-505.
3. Elia N, Culebras X, Mazza C, Schiffer E, Tramèr MR. Clonidine as an adjuvant to intrathecal local anesthetics for surgery: systematic review of randomized trials. *Reg Anesth Pain Med.* 2008;33:159-167.
4. Al-Ghanem SM, Massad IM, Al-Mustafa MM, Al-Zaben KR, Qudaisat IY, et al. Effect of Adding Dexmedetomidine versus Fentanyl to Intrathecal Bupivacaine on Spinal Block Characteristics in Gynecological Procedures: A Double Blind Controlled Study. *Am J Appl Sci.* 2009;(6):882-887.
5. Chaney MA. Side effects of intrathecal and epidural opioids. *Can J Anaesth.* 1995;42:891-903.
6. Al-Mustafa MM, Abu-Halaweh SA, Aloweidi AS, Murshidi MM, Ammari BA, et al. Effect of dexmedetomidine added

- to spinal bupivacaine for urological procedures. *Saudi Med J.* 2009;30:365-370.
7. Kanazi GE, Aouad MT, Jabbour-Khoury SI, Al Jassar MD, Alameddine MM, et al. Effect of low-dose dexmedetomidine or clonidine on the characteristics of bupivacaine spinal block. *Acta Anaesthesiol Scand.* 2006;50: 222-7.
8. Coursin D B, Coursin D B, Maccioli GA. Dexmedetomidine. *Current Opinion Critical Care.* 2001;7:221-226
9. Asano T, Dohi S, Ohta S, Shimonaka H, Iida H. Antinociception by epidural and systemic alpha 2 adrenoreceptor agonists and their binding affinity in rat spinal cord and brain. *AnesthAnalg.* 2000;90:400-7.
10. Mohamed AA, Fares KM, Mohamed SA. Efficacy of Intrathecally Administered Dexmedetomidine Versus Dexmedetomidine With Fentanyl in Patients Undergoing Major Abdominal Cancer Surgery. *Pain Physiol.* 2012;15:339-348.
11. Lawhead RG, Blaxall HS, Bylund BD. Alpha-2A is the predominant α-2 adrenergic receptor subtype in human spinal cord. *Anesthesiology.* 1992;77:983-91.
12. Ishii H, Kohno T, Yamakura T, Ikoma M, Baba H () Action of dexmedetomidine on the substantiagelatinosa neurons of the rat spinal cord. *Eur J Neurosci.* 2008;27:3182-3190.
13. Eisanach JC, De Kock M, Klimscha W. α-2 adrenergic agonists for regional anesthesia. *Anesthesiology.* 1996;85: 655-74.
14. Grewal A. Dexmedetomidine: New avenues. *J Anaesthesiol Clin Pharmacol.* 2011;27: 297-302.
15. Shimode N, Fukuoka T, Tanimoto M, Tashiro C, Tokunaga A, et al. The effects of dexmedetomidine and halothane on Fos expression in the spinal dorsal horn using a rat postoperative pain model. *Neurosci Lett.* 2003;343: 45-48.
16. Brummett CM, Norat MA, Palmisano JM, Lydic R. Perineural administration of dexmedetomidine in combination with bupivacaine enhances sensory and motor blockade in sciatic nerve block without inducing neurotoxicity in rat. *Anesthesiology.* 2008;109(3):502-11
17. Asano T, Dohi S, Ohta S, Shimonaka H, Iida H. Antinociception by epidural and systemic alpha 2 adreno receptor agonists and their binding affinity in rat spinal cord and brain. *Anesth Analg.* 2000; 90: 400—7.

**How to cite this article:** Chaudhry G, Bansal P, Singla D, Dureja J, Dahiya S. Comparison of Two Different Doses of Dexmedetomidine as an Adjuvant to Intrathecal Bupivacaine in Patients Undergoing Femur Surgery. *Ann. Int. Med. Den. Res.* 2016;2(1):94-7.

**Source of Support:** Nil, **Conflict of Interest:** None declared