

Clinical Profile of Patients with Tuberculous Pleural Effusion.

Sayed Khayyum Ali¹, Supriya Adiody²

¹Professor, Department of Tuberculosis and Respiratory Diseases, Malla Reddy Institute of Medical Sciences, Hyderabad, Telangana, India.

²Professor, Department of Tuberculosis and Respiratory Diseases, Jubilee Mission Medical College, Thrissur, Kerala, India.

ABSTRACT

Background: The diagnosis of tuberculous pleural effusion is by demonstration of the causative organism that is mycobacterium tuberculosis. Biochemical investigations like pleural fluid ADA which is shed by the lymphocytes is very non-specific. It is raised in other conditions like empyema, lymphoreticular malignancies, bronchogenic carcinoma, rheumatoid arthritis and tuberculosis (TB). The aim is to study the clinical profile of patients with tuberculous pleural effusion. **Methods:** A study was conducted on 60 patients at a public hospital. The patients were asked a detailed history with regards to chest pain and its nature, dyspnoea, cough, fever and constitutional symptoms. Also detailed past history about pleural tuberculosis and pleural aspiration/ICD insertion and AKT history was taken. Patients were then subjected to a clinical examination, which included a thorough general examination and detailed respiratory system examination in terms of decrease in movements of the chest on any particular side, bulge on any particular side, stony dull note on percussion, presence of shifting dullness, and decrease in breath sounds with decrease in vocal resonance. Patients were later subjected to an X ray examination for confirmation of the diagnosis. In some patients ultrasonography of the chest was done especially in loculated effusion for point marking. After confirming the diagnosis, patients were subjected to a transthoracic pleural aspiration using all aseptic precautions. **Results:** Most of the patients belonged to age group between 15-45 years i.e. 47 (78.33%). Thus the incidence in elderly age group was found to be less (21.67%). The symptom analysis of 60 patients revealed that almost all these patients had chest pain as predominant symptom (98%) followed by dry cough and dyspnoea on exertion, 98% and 88% respectively. The constitutional symptom such as loss of appetite was present in 93% of patients and weight loss in 85% of patients. The average duration of symptoms was 56.57 days. Right sided pleural effusion was more common than left side. There were no cases of bilateral pleural effusions at any given point of time during study period. **Conclusion:** Most common presentation of tuberculous pleural effusion was chest pain. Right sided pleural effusion was more common than left side.

Key words: Clinical profile, Tuberculosis, Pleural effusion.

INTRODUCTION

The diagnosis of tuberculous pleural effusion is by demonstration of the causative organism that is mycobacterium tuberculosis. Biochemical investigations like pleural fluid ADA which is shed by the lymphocytes is very non specific. It is raised in other conditions like empyema, lymphoreticular malignancies, bronchogenic carcinoma, rheumatoid arthritis and tuberculosis (TB).^[1,2]

Pleural biopsy can give the diagnosis but is an invasive and painful procedure with the risk of pneumothorax. When reviewing data regarding yield of pleural biopsy, it is important to distinguish histopathologic demonstration of granuloma from growth of M. tuberculosis in a culture of biopsy, though either is generally accepted as diagnostic. Certain clinical conditions like TB, plague, fungal infections, rheumatoid arthritis, and sarcoidosis may cause difficulty in distinguishing them histopathologically because of certain common feature.

Pleural fluid acid fast bacilli (AFB) smear is positive in less than 10% of cases. This phenomenon has been attributed to low bacillary counts in the fluid and the diminished likelihood of recovery of AFB in the absence of apparent parenchymal disease. Also the tuberculin test is negative in about 30% of patients because of the aggregation of T lymphocytes at the site of pleural focus. Diagnostic tests like PCR are too expensive and unaffordable for the common man. Hence it becomes very important to use less expensive and reliable tests like culturing the mycobacteria. Culture yield of mycobacteria from pleural fluid is less than 30%.^[3]

MATERIALS AND METHODS

A study was conducted on 60 patients at a public hospital. The patients were asked a detailed history with regards to chest pain and its nature, dyspnoea, cough, fever and constitutional symptoms. Also detailed past history about pleural tuberculosis and pleural aspiration / ICD insertion and AKT history was taken. Patients were then subjected to a clinical examination, which included a thorough general examination and detailed respiratory system examination in terms of decrease in movements of the chest on any particular side, bulge on any particular side, stony dull note on percussion,

Name & Address of Corresponding Author

Dr. Sayed Khayyum Ali
Professor and HOD,
Department of Tuberculosis and Respiratory Diseases,
Malla Reddy Institute of Medical Sciences,
Hyderabad, Telangana, India.
Email: sayedshafi@gmail.com

presence of shifting dullness, and decrease in breath sounds with decrease in vocal resonance. Patients were later subjected to an X ray examination for confirmation of the diagnosis. In some patients ultrasonography of the chest was done especially in loculated effusion for point marking.

Inclusion criteria:

All clinically suspected tuberculous pleural effusion having exudative character by Light's criteria and which on gram stain examination show no organisms

Exclusion criteria:

Those patients having parenchymal lesions on X ray chest

Those patients having hydro pneumothorax.

RESULTS

Most of the patients belonged to age group between 15-45 years i.e. 47 (78.33%). Thus the incidence in elderly age group was found to be less (21.67%) [Table 1]. The symptom analysis of 60 patients revealed that almost all these patients had chest pain as predominant symptom (98%) followed by dry cough and dyspnoea on exertion, 98% and 88% respectively. The constitutional symptom such as loss of appetite was present in 93% of patients and weight loss in 85% of patients [Table 2].

Table 1: Age and sex distribution of patients

Age group (years)	Male	Female	Total
15-25	14	6	20
25-35	13	3	16
35-45	8	3	11
45-55	2	3	5
55-65	2	1	3
65-75	4	1	5
Total	43 (71.66%)	17 (28.33%)	60 (100%)

Table 2: Distribution of study subjects as per their symptoms

Symptoms	Number	%
Chest pain	58	98
Dry cough	58	98
Dyspnoea	50	88
Fever	56	93
Loss of appetite	56	93
Loss of weight	51	85

Table 3: Distribution of study subjects as per their average duration of symptoms

Symptoms	Mean duration (days)
Chest pain	75.206
Breathlessness	56.57
Dry cough	70.327
Fever	63.79

The average duration of symptoms was 56.57 days. Right sided pleural effusion was more common than left side. There were no cases of bilateral pleural effusions at any given point of time during study period.

DISCUSSION

Pleural TB is a pauci bacillary infection. The diagnosis of pleural TB depends on the clinical history and pleural fluid characters such as lymphocytic predominant exudative pleural effusion with sugar low but, usually more than 40 mg/dl and ADA above cut off 47-53 units/L.^[2]

The definitive diagnosis of pleural TB still rests with demonstration of causative organisms either on direct microscopy or in the culture of the pleural fluid. The culture of pleural biopsy specimen from untreated tuberculosis pleural effusions has yield of 80%. Culture of pleural fluid, biopsy culture and

histopathology of pleural biopsy give a yield of 90-97%.^[1,2]

Certain clinical conditions like TB, plague, fungal infections, rheumatoid arthritis, and sarcoidosis may cause difficulty in distinguishing them histopathologically because of certain common features. The combined yield of direct acid fast smears of pleural fluid and pleural biopsy tissue along with culture of fluid and tissue exceeds 90%.^[4,5]

In this study, the incidence of pleural effusion is seen in young individuals. Mean age of presentation was 15-45 years. The incidence declines after age advances. This is what has been in contrast to western countries where the incidence increases after age advances because of low endemicity leading to infection at later age of life. The symptom analysis has revealed that predominant symptoms are pleuritic chest pain

(98%), loss of appetite (93%), followed by dyspnoea (88%) and fever (85%).

Our study has shown that right sided effusions were a little more as compared to left sided effusions. There were no cases of bilateral effusions seen.

REFERENCES

1. Fadda S. Recovery and susceptibility testing of mycobacterium tuberculosis from extra pulmonary specimens by the BACTEC radiometric method. J Clin Microbiol 1984;19:720-1.
2. Kamholz S. Pleural Tuberculosis. In: Tuberculosis. Rom WN, Garay SM editors. 1st ed. Little Brown and company. 1996. pp: 483-91.
3. Nagesh BS. Evaluation of polymerase chain reaction of mycobacterium tuberculosis in pleural fluid. Chest 2001;119:1737-41.
4. Pope RH. Sensitivity, specificity and predictive values of closed pleural biopsy. Arch Internal Med 1984;44:325-8.
5. Wali JP. Tuberculosis pleural effusion. In: Tuberculosis. Sharma SK editor. 1st ed. 2001 pp: 169-81.

How to cite this article: Ali SK, Adiody S. Clinical Profile of Patients with Tuberculous Pleural Effusion. Ann. Int. Med. Den. Res. 2016;2(1):193-95.

Source of Support: Nil, **Conflict of Interest:** None declared.