



## Ultra Sonographic Prostate Volume Measurement of Patients Over 40 Years in Age: A Single Center Study

Towhida Naheen<sup>1\*</sup>, Mizanur Rahman<sup>2</sup>, Kazi Md. Shahidur Rahman<sup>3</sup>, Nazmun Nahar<sup>4</sup>,  
Mohammed Iqbal<sup>5</sup>

<sup>1</sup>Department of Anatomy, Chattogram International Medical College (CIMC), Chattogram, Bangladesh.

Email: towhida.naheen@yahoo.com,

Orcid ID: 0000-0002-3566-5382

<sup>2</sup>Associate Professor, Department of Urology, Chattogram Medical College Hospital, Chattogram, Bangladesh.

Email: mizanuro@yahoo.com,

Orcid ID: 0000-0001-9966-2482

<sup>3</sup>Associate Professor, Department of Pathology, BGC Trust Medical College, Chandanaish, Chattogram, Bangladesh.

Email: shahid084@gmail.com,

Orcid ID: 0000-0002-6493-334X

<sup>4</sup>Associate Professor, Department of Pathology, Medical College for Women, Dhaka, Bangladesh.

Email: nahar085081@gmail.com,

Orcid ID: 0000-0002-6225-145X

<sup>5</sup>Associate Professor, Department of Anatomy, Monno Medical College, Manikganj, Bangladesh,

Email: iqbal085nahar082@gmail.com,

Orcid ID: 0000-0002-2346-361X

\*Corresponding author

Received: 17 May 2022

Revised: 23 July 2022

Accepted: 02 August 2022

Published: 22 August 2022

### Abstract

**Background:** Benign prostatic hyperplasia (BPH), also called prostate enlargement, is a noncancerous increase in size of the prostate gland. Symptoms may include frequent urination, trouble starting to urinate, weak stream, inability to urinate, or loss of bladder control. The objective of the study was to evaluate the prostate volume measurement for the Bangladeshi population over the age of 40 years by ultrasonography. **Material & Methods:** This prospective, observational study was conducted in the Department of Anatomy, Chattogram International Medical College (CIMC), Chattogram, Bangladesh during the period from January 2019 to December 2020. In total 157 suspected patients of benign prostatic hyperplasia were selected as the study population. All patients were clinically diagnosed for BPH, based on the present prostate symptoms and digital rectal examination. To measure the prostate volume, abdominal ultrasonography was performed for all the patients. After enucleation, another ultrasonogram was performed for all the patients to measure the existing sizes of the prostates of the patients. All the data were processed, analyzed, and disseminated by MS-word and SPSS programs as per need. **Results:** Finally, in this study in analyzing the volumes of the prostates of the participants according to the abdominal ultrasonographic reports of pre-operative stage we observed, in 9%, 34%, 31%, 30%, 21% and 32% patients, the prostate sizes (In cc) were <20, 21-40, 41-60, 61-80, 81-100 and >100 cc respectively. On the other hand, after enucleation, in 11.46%, 24.20%, 28.66%, 27.39%, 7.01% and 1.27% patients, the prostate sizes (In cc) were found <20, 21-40, 41-60, 61-80, 81-100 and >100 cc respectively. The mean changes of prostate sizes between pre- and post-operative stages among the participant was not significant where the P value was found 0.464. **Conclusion:** The findings of this study support the applications of abdominal ultrasonographic evaluation for suspected benign prostatic hyperplasia patients to know about the exact volumes of their prostates for selecting the appropriate surgical approach.

**Keywords:-** Prostate gland, Volume, Benign prostatic hyperplasia, BPH, Ultrasonography.

### INTRODUCTION

Benign prostatic hyperplasia is a histologic diagnosis status characterized by proliferation

of the 'cellular elements' of the prostate, which may lead to an enlarged prostate gland. Ultrasonography is a very widely used method to determine prostate volume or size. Benign

prostatic hyperplasia (BPH), also known as prostate gland enlargement is a very common surgical problem accounting for 20% of elective admissions in the surgical wards.<sup>[1]</sup> Although BPH is diagnosable by clinical examinations, 'information about gland volume/size', presence of nodules, and 'calcification' are important for selecting the proper management.<sup>1</sup> The decision on whether the patient requires surgery as well as the selection of the best surgical method is dependent upon the size of the prostate.<sup>[2]</sup> Ultrasonography is considered the most prominent method to determine prostate volume or size. Ultrasonography, specifically transrectal ultrasonography (TRUS), is the most commonly applied tool to estimate prostate volume.<sup>[3]</sup> The ultrasonography computes the 'total volume of the prostate by measuring the length, height as well as the width of the gland and multiplying the product by a coefficient of  $\pi/6$  (0.52), which is also known as the prolate ellipsoid formula.<sup>[4]</sup> To estimate the prostate volume, the same formula can also be used in MRI (Magnetic Resonance Imaging) modalities.<sup>[4]</sup> Besides these, another geometrical model is known as bullet formula ( $L \times H \times W \times 5\pi/24$ ) or ( $L \times H \times W \times 0.65$ ) was introduced in 2009 as a potentially superior formula for estimating prostatic volume.<sup>[5]</sup> In this study, our focus was on ultra-sonographic findings.

## Objectives

### General objective

Evaluation of the measurement of prostate volume by Ultrasonography over the age of 40 years Patients.

### Specific objectives

- To evaluate the demographic profile of the patient.
- To assess the age status of the patients.
- To assess the symptoms among the patients.

## MATERIAL AND METHODS

This prospective, observational study was conducted in the Department of Anatomy, Chattogram International Medical College (CIMC), Chattogram, Bangladesh during the period from January 2019 to December 2020. In total 157 suspected patients of benign prostatic hyperplasia were selected as the study population. Ethical committee of the hospital approved this study. According to the exclusion criteria of this study, patients who had the history of pelvic or prostatic surgery, acute/chronic urinary retention, recurrent urinary tract infection (UTI) or bladder stones, cases with chronic or acute prostatitis within the previous 3 months and known cases of prostatic carcinoma, hypertension and diabetes mellitus (DM) were excluded from our study. All patients were clinically diagnosed for BPH, based on the present prostate symptoms and digital rectal examination. By using the transducer in a 45° C caudally angulated position, preoperative abdominal ultrasound examinations were performed with a full bladder (200-300 ml urine). To measure the prostate volume, abdominal ultrasonography was performed for all the patients. A 3.5 MHz transducer was applied to measure the width, height, and length of the prostate. After enucleation, another ultrasonogram was performed for all the patients to measure the existing sizes of the prostates of the patients. A pre-designed questioner has used inpatient

data collection. All the data were processed, analyzed, and disseminated by MS-word and SPSS programs as per need.

## RESULTS

In this study, among total 157 suspected patients of benign prostatic hyperplasia were finally enrolled as the study population. To measure the prostate volume abdominal ultrasonography was performed for all the patients. By using the transducer in a 45° C caudally angulated position, preoperative abdominal ultrasound examinations were performed with a full bladder (200-300 ml urine). A 3.5 MHz transducer was applied to measure the width, height, and length of the prostate. After enucleation, another ultrasonogram was performed for all the patients to measure the existing sizes of the prostates of the patients. The mean ( $\pm$ SD) age of the participants was  $62.20 \pm 5.14$  years. The highest number of participants were found

from the 61-70 year's age group which was 38%. Besides this, 24% of participants were from 51-60 years, 21% were from >70 and the rest 17% were from 41-50 year's age groups. In analyzing the signs and symptoms among the patients we found urgency, dribbling, hesitancy, acute retention, dysuria, and palpable bladder in 98%, 96%, 91%, 85%, 81% and 74% participants respectively. In this study, in analyzing the volumes of the prostates of the participants according to the abdominal ultra-sonographic reports of pre-operative stage we observed, in 9%, 34%, 31%, 30%, 21% and 32% patients, the prostate sizes (In cc) were < 20, 21-40, 41-60, 61-80, 81-100 and >100 cc respectively. On the other hand, after enucleation, in 11.46%, 24.20%, 28.66%, 27.39%, 7.01% and 1.27% patients, the prostate sizes (In cc) were found < 20, 21-40, 41-60, 61-80, 81-100 and >100 cc respectively. The mean changes of prostate sizes between pre- and post-operative stages among the participant was not significant where the P value was found 0.464.

**Table 1:** Age Group Distribution Among the Participants (N=157)


Age group	n	%
41-50 yrs.	27	17.0
51-60 yrs.	37	24.0
61-70 yrs.	60	38.0
> 70 yrs.	33	21.0

**Table 2:** Signs and Symptoms Among the Participants (N=157)


Symptoms	n	%
Urgency	154	98.0
Dribbling	150	96.0
Hesitancy	143	91.0
Acute retention	134	85.0
Dysuria	127	81.0
Palpable bladder	116	74.0

**Table 3:** Pre- and post-operative prostatic volumes measured by abdominal ultrasound (N=157)


Size (cc)	Preoperative		After Enucleation		P Value
	n	%	n	%	
< 20 cc	9	5.73	18	11.46	0.464
21-40 cc	34	21.66	38	24.20	
41-60 cc	31	19.75	45	28.66	
61-80 cc	30	19.11	43	27.39	
81-100 cc	21	13.38	11	7.01	
>100 cc	32	20.38	2	1.27	




**Figure 1:** Age distribution of the participants (N=157)




**Figure 3:** Pre- and post-operative prostatic volumes measurement of the participants (N=157)



**Figure 2:** Sign & Symptoms of the participants (N=157)





## DISCUSSION

The aim of the present study was to evaluate the prostate volume measurement for the Bangladeshi population over the age of 40 years by ultrasonography. The highest number of participants were found from the 61-70 year's age group which was 38%. Besides this, 24% of participants were from 51-60 years, 21% were from >70 and the rest 17% were from 41-50 year's age groups. In our study, in analyzing the volumes of the prostates of the participants according to the abdominal ultra-sonographic reports of pre-operative stage we observed, in 9%, 34%, 31%, 30%, 21% and 32% patients, the prostate sizes (In cc) were 100 cc respectively. Assessment of the size of the prostate is an important factor in the management of Benign prostatic hyperplasia (BPH).<sup>[6]</sup> In such cases abdominal ultrasonography is in need. Intravenous urography or IVU and cystoscopy are not useful at all for determining the exact prostatic size or selecting the proper surgical approach.<sup>[7]</sup> As per the recommendation of Roehrborn et al., the best predictor of prostatic weight is abdominal ultrasound ( $r = 0.975$ ).<sup>[8]</sup>

The reliability of abdominal ultrasonography in the measurement of the volumes of the prostate was also confirmed by many other studies.<sup>[9,10]</sup> In this current study, after enucleation, in 11.46%, 24.20%, 28.66%, 27.39%, 7.01% and 1.27% patients, the prostate sizes (In cc) were found 100 cc respectively. The majority of the enucleated glands weighed less than 60 gm (71.7%) suggests the procedure of choice would have been transurethral resection of the prostate (TURP).<sup>[11]</sup> Surgeons can depend upon the findings of ultrasonography in selecting the appropriate treatment method for BPH patients. The correlation coefficient in several series generally shows a statistically significant correlation between prostatic volume measured on abdominal ultrasonography and the weight of the prostate removed at surgery, indicating that it is a useful tool for selecting the appropriate surgical approach. Some studies have shown a similar correlation. Styles et al. found that abdominal estimation of prostatic volume, correlated well with the 'transrectal method', and good inter-observer agreement was found with the use of both modalities.<sup>[12]</sup> Ishida et al. demonstrated that 'abdominal measurement of prostatic volume' correlated well with the resected weight ( $r=0.956$ ).<sup>[13,14]</sup>

## CONCLUSIONS

The findings of this study support the applications of abdominal ultrasonographic evaluation for suspected benign prostatic hyperplasia patients to know about the exact volumes of their prostates for selecting the appropriate surgical approach. For getting more specific information regarding this issue we would like to recommend conducting more studies in several places with larger sample sizes.

## Limitations of the Study

This was a single-centered study with a small sample. Due to the limited sample size, the

findings of this study might not reflect the exact scenario of the whole country.

## REFERENCES

1. Getahun GM, Kebede AG. Comparison of prostatic volume measured with abdominal ultrasound and prostatic weight determined after open enucleation performed in Gondar University Hospital, Ethiopia. *Afr J Urol.* 2015;14(2), 86-89.
2. Watanabe T, Miyagawa I. New simple method of transabdominal ultrasound to assess the degree of benign prostatic obstruction: Size and horizontal shape of the prostate. *Int J Urol.* 2002;9(4):204-9.
3. Harvey CJ, Pilcher J, Richenberg J, Patel U, Frauscher F. Applications of transrectal ultrasound in prostate cancer. *Br J Radiol.* 2012;85(1): S3-17.
4. Lee JS, Chung BH. Transrectal ultrasound versus magnetic resonance imaging in the estimation of prostate volume as compared with radical prostatectomy specimens. *Urol Int.* 2007;78(4):323-7.
5. MacMahon PJ, Kennedy AM, Murphy DT, Maher M, McNicholas MM. Modified prostate volume algorithm improves transrectal US volume estimation in men presenting for prostate brachytherapy. *Radiology.* 2009;250(1):273-80.
6. Prassopoulos P, Charoulakis N, Anezinis P, Daskalopoulos G, Cranidis A, Gourtsoyiannis N. Suprapubic versus Tran's rectal ultrasonography in assessing the volume of the prostate and the transition zone in patients with benign prostatic hyperplasia. *Abdom Imaging.* 1996;21(1):75-7.
7. Thumann RC Jr. Estimation of the weight of the hyperplastic prostate from the cyst urethrogram. *Am J Roentgenol Radium Ther Nucl Med.* 1951;65(4):593-4.
8. Roehrborn CG, Chinn HK, Fulgham PF, Simpkins KL, Peters PC. The role of transabdominal ultrasound in the preoperative evaluation of patients with benign prostatic hypertrophy. *J Urol.* 1986; 135(6):1190-3.
9. Rathaus V, Richter S, Nissenkorn I, Goldberg E. Transperineal ultrasound examination in the evaluation of prostatic size. *Clin Radiol.* 1991;44(6):383-5.
10. Bartsch G, Egender G, Hubscher H, Rohr H. Sonometrics of the prostate. *J. Urol.* 1982;127(6):1119-21.
11. Vilmann P, Hancke S, Strange Vognsen HH, Nielsen K, Sorensen SM. The reliability of transabdominal ultrasound scanning in the determination of prostatic volume. An autopsy study. *Scand J Urol Nephrol.* 1987;21(1):5-7.
12. Styles RA, Neal DE, and Powell PH. Reproducibility of measurement of prostatic volume by ultrasound. Comparison of transrectal and transabdominal methods. *Eur Urol.* 1988;14(4):266-9.
13. Smith HJ, Haveland H. Pre-operative and post-operative volume try of the prostate by transabdominal ultrasonography. *Br J Urol.* 1982; 54(5):531-5.
14. Ishida N, Tsurumaki O, Igarashi S, Maezawa H, Yamamoto R, Hosoi Y, et al. [The evaluation of simple estimation method of prostate size by transabdominal ultrasound]. *Nippon Hinyokika Gakkai Zasshi.* 1989;80(6):832-7.

Source of Support: Nil, Conflict of Interest: None declared